



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Current Topic / Актуелна тема

Nikola Milošević^{1,*}, Suzana Stojanović-Rundić^{1,2}, Srđan Milanović¹

Pancreatic carcinoma – diagnosis and modern multimodal treatment

Карцином панкреаса – дијагноза и савремено мултимодалног лечења

¹Institute for Oncology and Radiology of Serbia, Belgrade, Serbia;

²University of Belgrade, Faculty of Medicine, Belgrade, Serbia

Received: April 23, 2024

Revised: August 18, 2024

Accepted: September 1, 2024

Online First: September 3, 2024

DOI: <https://doi.org/10.2298/SARH240423071M>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

* **Correspondence to:**

Nikola MILOŠEVIĆ

Institute for Oncology and Radiology of Serbia, Pasterova 14, 11000 Belgrade, Serbia

E-mail: milosevicnikola43@gmail.com

Pancreatic carcinoma – diagnosis and modern multimodal treatment

Карцином панкреаса – дијагноза и савремено мултимодалног лечења

SUMMARY

Pancreatic cancer is one of the most aggressive tumors and is among the top ten most common malignancies in the world. This is a disease of older adults, and men get it more often. Pancreatic carcinomas risk factors are obesity and type II diabetes, smoking and alcohol. Symptoms of the disease include obstructive jaundice, loss of appetite, fatigue and back pain. The diagnosis of pancreatic cancer involves computed tomography of the thorax, abdomen and pelvis or magnetic resonance imaging of the abdomen and pelvis, and endoscopic ultrasound with biopsy. The most common histological type of pancreatic cancer is ductal adenocarcinoma. The TNM classification is used to determine the stage of the disease. Pancreatic cancer treatment is complex, multidisciplinary and multimodal, and involves the use of surgery, chemotherapy and radiotherapy, alone or in different combinations. Surgery is the main treatment modality for these tumors, especially in localised stages. Chemotherapy is applied in all forms in the treatment of pancreatic cancer as neoadjuvant, adjuvant and systemic. Immunotherapy, as the newest type of treatment, is used in a limited way in the metastatic phase of pancreatic cancer. The role of radiotherapy in the treatment of pancreatic cancer is still debated, and it is most often applied in a neoadjuvant and palliative approach. Palliative therapy and care are an indispensable part of the treatment of patients with pancreatic cancer.

Keywords: pancreatic cancer; treatment; surgery; chemotherapy; radiotherapy

САЖЕТАК

Карцином панкреаса је један од најагресивнијих тумора и налази се међу првих десет најчешћих малигнитета у свету. Ово је болест старијих особа, а мушкарци оболевају чешће. Карциноми панкреаса су тумори са утицајем генетских наследних синдрома и стечених фактора као што су гојазност и дијабетес тип II, пушење и алкохол. Симптоми болести укључују опструктивни иктерус, губитак апетита, умор, губитак тежине, инсуфицијенцију панкреаса, бол у трбуху, бол у леђима, и умор. Дијагноза карцинома панкреаса подразумева ођређивање тумор маркера и биомаркера, компјутеризована томографија (ЦТ) торакаса, ЦТ / магнетна резонанца абдомена и мале карлице, ендоскопски ултразвук са биопсијом. Најчешћи хистолошки тип карцинома панкреаса је дуктални аденокарцином. За одређивање стадијума болести се користи ТНМ класификација. Лечење карцинома панкреаса је мултидисциплинарно и мултимодално, а подразумева примену хирургије, хемотерапије и радиотерапије, самостално или у различитим комбинацијама. Хирургија је основни модалитет лечења ових тумора, нарочито у раном стадијуму. Хемотерапија се у лечењу карцинома панкреаса примењује као неoadјувантна, адјувантна и системска. Имунотерапија се као најновији вид лечења ограничено примењује у метастаској фази карцинома панкреаса. Улога радиотерапије у лечењу карцинома панкреаса је врло контраверзна, о њој се још увек дебатује, а најчешће се примењује у неoadјувантном и палијативном приступу. Палијативна терапија и нега су незаобилазни део током лечења пацијената са карциномом панкреаса.

Кључне речи: карцином панкреаса; лечење; хирургија; хемотерапије; радиотерапија

INTRODUCTION

Pancreatic carcinomas are considered one of the deadliest malignancies because they are usually in advanced stage at the time of diagnosis.

EPIDEMIOLOGY AND ETIOLOGY

Pancreatic cancer is among the top ten cancers in the world in terms of incidence and the seventh most common cause of dying from cancer [1]. In the Republic of Serbia, pancreatic cancer is the sixth most common cancer.

Pancreatic cancer is a multifactorial disease. Genetic factors include a positive family history. Demographic risk factors, age and sex, indicate that the disease is more common over the age of 50 and with a male predominance. Acquired risk factors include obesity, type II diabetes, smoking and alcohol [2].

SYMPTOMS AND DIAGNOSIS

Early-stage pancreatic cancer rarely causes symptoms, and when symptoms develop, it is usually at an advanced stage. Symptoms includes obstructive jaundice, loss of appetite, fatigue, weight loss, and abdominal or back pain.

At the time of diagnosis, about 50% of patients have metastatic disease, about 35% have locally advanced disease, and 15% have local disease [3].

Diagnostics of pancreatic cancer includes computerized tomography (CT) of the thorax, magnetic resonance (MR) or CT of the abdomen and pelvis, MR cholangiopancreatography (MRCP), endoscopic ultrasound (EUS) with biopsy and pathohistological analyses.

CT is widely accepted in the diagnosis of pancreatic cancer. With modern multiphase and multidetector high-resolution CT with multiplanar reconstructions and contrast, the sensitivity of pancreatic cancer detection is 89%, and the specificity is 90% [4].

MR of the abdomen and pelvis provides a much better image of the soft tissues, especially in cases where the CT findings are inconclusive. MR with using diffusion sequences (DWI), the sensitivity increases to 92-96%, and the specificity to 97-99% [4].

EUS is not used as a routine method in the detection of pancreatic cancer, but for the detection of small tumors that are not visible on CT and are suspicious on MR. EUS is primarily used to biopsy suspicious changes in the head of the pancreas to obtain PH findings. The sensitivity of this method is 89–91% and the specificity is 81–86% [5].

Pancreatic carcinomas form a very heterogeneous group of tumors and ductal adenocarcinomas cancers make about 90% of all pancreatic [6]. Although histopathology remains the most reliable method for establishing a definitive diagnosis, it may be difficult to distinguish pancreatic carcinomas from metastases without immunohistochemistry (IHH) [7]. Progress in molecular biology and pathology have led to more precise diagnosis, subtyping, tumor classification and prediction of therapeutic response. Pancreatic cancer genome is very heterogeneous and includes gene mutations and abnormalities (BRCA1, BRCA2, TP53, KRAS, etc.), microsatellite instability (MSI) as well as impaired expression of growth factors (HER2) [8].

STAGING

The TNM classification is used to present the stage of pancreatic cancer [9]. The lymphovascular and perineural invasion of pancreatic carcinoma is prominent. The most common sites of pancreatic cancer metastasis are the liver (90%), lymph nodes and lungs (25%), peritoneum (20%) and bones (10-15%) [10].

TREATMENT

Five-year survival of a patient with pancreatic cancer is very modest, about 30% in the local stage, about 10% in the locally advanced stage, and 5% in the metastatic stage of the disease [8]. Pancreatic cancer treatment is multidisciplinary and multimodal. The basic type of treatment for early pancreatic cancer involves surgery, for locally advanced cancers a

combination of chemotherapy with/or without radiotherapy and surgery, and for metastatic cancers, chemotherapy and radiotherapy. The most widely used recommendations in the treatment of pancreatic cancer are the National comprehensive cancer network (NCCN) [11] and the European society of medical oncology (ESMO) [12].

SURGERY

The primary approach in the treatment of pancreatic cancer is radical surgery and involves complete removal of the visible tumor and obtaining histologically free margins, i.e. microscopically negative margin means a distance of less than 1mm from cancer cells.

In terms of resectability, pancreatic cancers can be estimated as resectable, borderline resectable or unresectable [11].

Treatment is surgical for resectable tumors. Borderline resectable and locally advanced tumors are operated upon after neoadjuvant therapy and conversion to a resectable stage. Radical surgical procedures are pancreaticoduodenectomy (Kausch-Whipple procedure), distal pancreatectomy and total pancreatectomy.

Pancreaticoduodenectomy (Kausch-Whipple procedure) is an operation for cancer of the head of the pancreas that includes the removal of the distal part of the stomach, duodenum, the end of the common bile duct, the gallbladder and the head of the pancreas with at least a 1 cm margin from the tumor. Reconstruction can be achieved in the form of pancreaticjejunostomy or pancreaticogastrostomy [13]. Lymphadenectomy involves dissection of the associated lymph nodes and standard dissection includes the infra- and suprapyloric lymph nodes, along the common hepatic artery, lymph nodes in the hepaticoduodenal ligament, around the pancreatic head and around the proximal part of the superior mesenteric artery and superior mesenteric vein. Distal pancreatectomy is an operation for carcinoma of the body and tail of the pancreas with "en-block" splenectomy, and involves mobilization of the body and tail of

the pancreas, ligation of the splenic artery and splenic vein, lymphadenectomy, transection of the pancreas and ligation of the pancreatic duct. Total pancreatectomy is a combination of pancreaticoduodenectomy and distal pancreatectomy in patients with multilocular, large, and centrally located pancreatic tumors. It includes "en bloc" resection of the pancreas, gall bladder, duodenum and spleen with accompanying lymphadenectomy and reconstruction with hepaticojejunostomy and duodenojejunostomy.

The operation can be classic (open) and minimally invasive (laparoscopic and robotic). The minimally invasive techniques in the treatment of pancreatic cancer are not inferior comparing to open surgery in terms of surgical and oncological outcome. Laparoscopic technique may lead to a reduced hospital stay and postoperative complications and increased percentage of marginal clearance resections and harvested lymph nodes [14]. Robotic surgery is promising technique that may lead to shorter hospitalization and lower postoperative complications but longer operative time [14]. Minimally invasive technique should be performed by experienced surgeons in high-volume centers and randomized trials with a large number of patients are necessary to examine the benefits of new technique, especially robotic surgery.

CHEMOTHERAPY

In the treatment of pancreatic cancer, chemotherapy (HT) is applied in almost all stages of the disease (except in early stages), as preoperative (neoadjuvant), postoperative (adjuvant) and systemic (palliative).

The role of neoadjuvant HT is manifested in marginally resectable pancreatic cancers because it significantly improves overall survival (OS) compared to "up-front" surgery (median OS 19 months vs. 29 months with neoadjuvant HT) [15], usually in combination with radiotherapy. In locally advanced unresectable pancreatic cancer with median OS of about 11-

14 months [16], neoadjuvant HT is essentially induction, with the aim of reducing the size of the tumor and regional lymph nodes, to facilitate resection rate (up to 28%) and increase the chance of achieving R0 resection (up to nearly 70%), thus prolonging OS [17]. It can be combined with radiotherapy. Neoadjuvant HT can be applied as polytherapy with 5-FU leucovorin, oxaliplatin and irinotecan (FOLFIRINOX) or as doublet therapy with gemcitabine-cisplatin/gemcitabine-nabpaclitaxel (GP/GN). FOLFIRINOX provides a slightly greater benefit to overall survival (median OS 33,4 months with FOLFIRINOX and surgery vs. 27,9 months with GN and surgery) [17], but also slightly more pronounced toxic effects, so it is used in patients with preserved performance (ECOG PS 0-1), and for more fragile patients gemcitabine-based HT is a good alternative.

Adjuvant HT is indicated in operated patients with pancreatic cancer regardless of resection margin status, in poor prognostic factors such as younger patients, poor tumor differentiation (gr 3-4), T3 and T4 disease stage and positive ln (N+). In patients with preserved ECOG PS, modified (m) FOLFIRINOX is used as standard. PRODIGE 24/CCTG PA6 trial is the most relevant study on the role of adjuvant HT and demonstrated significantly longer survival with (m) FOLFIRINOX (median OS 53,5 months vs. 35,5 months in group with gemcitabine) [18], which is much longer OS than those patient with only observation after resection (median OS is 22,3 months) [19]. A good alternative to standard HT is gemcitabine-cisplatin doublet, while mono-gemcitabine and 5FU/LV is reserved for ECOG PS 2-3 patients [12].

Systemic HT is applied in the metastatic phase of the disease and has a palliative purpose. The same types of cytostatics are used. In the first line, in good ECOG PS 0-1, FOLFIRINOX is preferred over GN (median OS 9,6 months with FOLFIRINOX vs. 6,1 months with GN) [20], which is not a significant increase of OS (median OS in metastatic pancreatic cancer is 4,6-8,1 months) [21]. In fragile patients, mono-gemcitabine or mono-capecitabine is given [11]

[12]. In the second line, in patients with ECOG PS 0-1, who were previously treated with FOLFIRINOX, GN can be used, while in patients who received gemcitabine-based HT in the first line, nanosomal irinotecan-5FU-LV is used. Other cytostatic combinations in the metastatic phase of the disease are GP, capecitabine-oxaliplatin (Cape-OX) and oxaliplatin-5FU-LV (OFF) [11] [12].

New developments in cytotoxic chemotherapy imply NALIRIFOX (combination of liposomal irinotecan, fluoracil, leucovorin and oxaliplatin) and it is possible option for frontline therapy in previously untreated patients with metastatic pancreatic cancer [22]. Some of ongoing trials include an efficiency assessment of combination of small molecule and cytotoxic drug in therapy of metastatic pancreatic cancer, like AVENGER 500 trial (mFOLFIRINOX and CPI-613) and NCT03126435 trial (endoTAG-1 and gemcitabine) [23].

IMMUNOTHERAPY

Immunotherapy in the treatment of pancreatic cancer is limited and for now is only used in metastatic disease. Poly ADP ribose polymerase enzyme inhibitors (PARP inhibitors) such as olaparib are used, followed by "checkpoint" inhibitors such as pembrolizumab (they act on receptors for PD-1 and PD-L1 protein in the tumor cell). Olaparib is administered as maintenance therapy in patients with metastatic pancreatic cancer harboring a BRCA gene mutation (about 4% of patients) after first-line platinum-based chemotherapy [24]. Pembrolizumab in unresectable pancreatic cancers can lead to a good partial remission of the disease [25].

CAR T-cell therapy is a promising development in immunotherapy of pancreatic cancer and is driving future clinical trials. Ongoing immunotherapy trials include combination of PARP inhibitors and PD-1 inhibitor (SWOG2001 trial) and PD-1 inhibitor and small molecule (SX-682) (NCT04477343 trial) [22].

RADIOTHERAPY

The role of radiation in the treatment of pancreatic cancer is still insufficiently defined. State-of-the-art radiotherapy techniques such as intensity modulated radiotherapy (IMRT), volumetric modulated arc therapy (VMAT) and stereotactic "body" radiotherapy (SBRT) are used [26]. 4D-CT planning and image-guided radiotherapy (IGRT) are recommended. However, there is still no consensus regarding the fractionation of radiotherapy in the treatment of pancreatic cancer. Radiotherapy (RT) in the treatment of pancreatic cancer can be preoperative (neoadjuvant), postoperative (adjuvant) and palliative, usually at the same time as chemotherapy to enhance the effect of the therapy.

The real role of neoadjuvant HRT is in marginally resectable pancreatic cancers because they are at increased risk of R1 resection, and RT can lead to downsizing and/or downstaging of the tumor, which increases the chance of achieving R0 resection. Landmark study on the role of neoadjuvant HRT is the PREOPANC study with 250 patients that compared the 5-year OS of two groups of patients, treated with surgery alone plus adjuvant HT or a combination of preoperative HRT plus surgery and adjuvant HT (HT with gemcitabine and RT with TD 36Gy in 15 fr.). The percentage of R0 resection was 72% vs. 43%, 5-year OS rate was 20,5% vs. 6,5% (HRT and surgery group vs. surgery group) [27]. Today, standard therapy includes induction HT for 2-3 months, FOLFIRINOX or GN, and then concomitant HRT with 5-FU or capecitabine or gemcitabine [12]. Radiotherapy doses of neoadjuvant HRT for borderline resectable pancreatic cancer are varied starting from standard fractionation (45-54Gy in 25-30 fractions), through hypofractionation (36Gy in 15 fractions, 30Gy in 10 fractions) to stereotaxy (28-30Gy in 5 fractions, 33-40Gy in 5 fractions) [28]. In locally advanced unresectable pancreatic cancer, neoadjuvant therapy can lead to conversion to potentially resectable cancer. It was demonstrated in the CONKO-007 trial with over 300 patients that compared the OS of two groups of patients, treated with induction HT and sequential HT or preoperative HRT plus

surgery (induction HT with gemcitabine/FOLFIRINOX and RT with TD 50,4Gy in 28 fr.). The percentage of R0 resection was 25% vs. 18%, pCR was 10% vs. 0% and 2-year OS rate was similar (HT and surgery group vs. HRT and surgery group) [29]. The optimal therapeutic approach in the treatment of locally advanced unresectable pancreatic cancer remains controversial, with conversion induction HT followed by neoadjuvant HRT being more commonly used in the US, while induction HT is favored in most European countries. Radiotherapy doses of neoadjuvant HRT of locally advanced unresectable pancreatic cancer are also different starting from standard fractionation (50.4-54Gy in 28-30 fractions) which is less frequently used, through hypofractionation (36Gy in 15 fractions, 30Gy in 10 fractions) and ablative RT (67.5Gy in 15 fractions and 75Gy in 25 fractions) to stereotaxy (30Gy in 5 fractions, 33-35Gy in 5 fractions and 45Gy in 6 fractions) [28].

Adjuvant HRT is not a therapeutic standard in operated patients with pancreatic cancer because the role of adjuvant RT in the modern era of new and more effective systemic therapies remains unclear [30]. Adjuvant HRT can be considered in pT3 stage and pN+ cases, and RT doses would be 50-55Gy in 25-30 fractions [28].

Palliative radiotherapy is carried out in order to control local symptoms and improve the quality of life of patients in an advanced stage of malignant disease. Indications are pain, bleeding, bone and brain metastases, and radiotherapy doses are 8Gy in 1 fraction, 16Gy in 4 fractions, 20Gy in 5 fractions or 30Gy in 10 fractions.

PALLIATIVE THERAPY AND CARE

Palliative therapy and care are very important parts of pancreatic cancer treatment. It encompasses drug therapy (non-opioid, opioid and co-analgetics), palliative surgical interventions or endoscopic placement of stents in case of obstruction (gastrojejunostomy or biliary-digestive anastomosis) and palliative RT for bone and brain metastases.

CONCLUSION

Pancreatic cancer is an aggressive invasive tumor with a modest 5-year survival. The treatment of these cancers is complex and challenging, and involves the use of surgery, chemotherapy and immunotherapy as well as radiotherapy, alone or in different combinations. Surgery is the main way of treating pancreatic cancer, especially in the early stages of the disease. Various chemotherapy regimens are indispensable in the treatment of pancreatic cancer in the neoadjuvant, adjuvant and systemic approach. Immunotherapy as the newest type of therapy is emerging in the treatment of selected cases of these cancers, especially in systemic disease. Radiotherapy has an unclear role in the treatment of these tumors, and it is used in marginally resectable and locally advanced cancers in a neoadjuvant approach, usually in combination with chemotherapy, but also in a systemic approach. Palliative therapy is complementary and important part of pancreatic cancer therapy.

Ethics: This article was written in accordance with the ethical standards of the journal and the institutions.

Conflict of interest: None declared.

REFERENCES

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021; 71 (3): 209–49. [DOI: 10.3322/caac.21660] [PMID: 33538338]
2. Ke T, Lophatananon A, Nieroda M, Kyriazis D, Manias G, Wajid U, et al. Risk Factors of Pancreatic Cancer: A Literature Review. *Cancer Rep Rev.* 2022; 6: 1–10. [DOI: 10.15761/CRR.1000241]
3. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. *CA Cancer J Clin.* 2021;71 (1): 7–33. [DOI: 10.3322/caac.21654] [PMID: 33433946]
4. Elbanna KY, Jang HJ, Kim TK. Imaging diagnosis and staging of pancreatic ductal adenocarcinoma: a comprehensive review. *Insights Imaging.* 2020; 11 (1): 58. [DOI: 10.1186/s13244-020-00861-y] [PMID: 32335790]
5. Rhee H, Park MS. The Role of Imaging in Current Treatment Strategies for Pancreatic Adenocarcinoma. *Korean J Radiol.* 2021; 22 (1): 23–40. [DOI: 10.3348/kjr.2019.0862] [PMID: 32901458]
6. Haerberle L, Esposito I. Pathology of pancreatic cancer. *Transl Gastroenterol Hepatol.* 2019; 4: 50. [DOI: 10.21037/tgh.2019.06.02] [PMID: 31304427]
7. Luu TT. Review of Immunohistochemistry Biomarkers in Pancreatic Cancer Diagnosis. *Front Oncol.* 2021; 11: 799025. [DOI: 10.3389/fonc.2021.799025] [PMID: 34988027]
8. Liu J, Mroczek M, Mach A, Stępień M, Aplas A, Pronobis-Szczylik B, et al. Genetics, Genomics and Emerging Molecular Therapies of Pancreatic Cancer. *Cancers (Basel).* 2023; 15 (3): 779. [DOI: 10.3390/cancers15030779] [PMID: 36765737]
9. Roalsø M, Aunan JR, Søreide K. Refined TNM-staging for pancreatic adenocarcinoma - Real progress or much ado about nothing? *Eur J Surg Oncol.* 2020;46(8):1554–7. [DOI: 10.1016/j.ejso.2020.02.014] [PMID: 32107094]
10. Wu L, Zhu L, Xu K, Zhou S, Zhou Y, Zhang T. Clinical significance of site-specific metastases in pancreatic cancer: a study based on both clinical trial and real-world data. *J Cancer.* 2021; 12 (6): 1715–21. [DOI: 10.7150/jca.50317] [PMID: 33613759]
11. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology. Pancreatic Adenocarcinoma. Version 2; 2024. https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf
12. Conroy T, Pfeiffer P, Vilgrain V, Lamarca A, Seufferlein T, O'Reilly EM, et al. ESMO Guidelines Committee. Pancreatic cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol.* 2023; 34 (11): 987–1002. [DOI: 10.1016/j.annonc.2023.08.009] [PMID: 37678671]
13. Mastalier B, Cauni V, Tihon C, Septimiu Petrutescu M, Ghita B, Popescu V, et al. Pancreaticogastrostomy versus Pancreaticojejunostomy and the Proposal of a New Postoperative Pancreatic Fistula Risk Score. *J Clin Med.* 2023; 12 (19): 6193. [DOI: 10.3390/jcm12196193] [PMID: 37834836]
14. Yan Y, Hua Y, Chang C, Zhu X, Sha Y, Wang B. Laparoscopic versus open pancreaticoduodenectomy for pancreatic and periampullary tumor: A meta-analysis of randomized controlled trials and non-randomized comparative studies. *Front Oncol.* 2023; 12: 1093395. [DOI: 10.3389/fonc.2022.1093395] [PMID: 36761416]
15. van Dam JL, Janssen QP, Besselink MG, Homs MYV, van Santvoort HC, van Tienhoven G, et al. Dutch Pancreatic Cancer Group. Neoadjuvant therapy or upfront surgery for resectable and borderline resectable pancreatic cancer: A meta-analysis of randomised controlled trials. *Eur J Cancer.* 2022; 160: 140–9. [DOI: 10.1016/j.ejca.2021.10.023] [PMID: 34838371]
16. Bratlie SO, Wennerblom J, Vilhav C, Persson J, Rangelova E. Resectable, borderline, and locally advanced pancreatic cancer-"the good, the bad, and the ugly" candidates for surgery? *J Gastrointest Oncol.* 2021; 12 (5): 2450–60. [DOI: 10.21037/jgo-2020-slapc-04] [PMID: 34790406]
17. Eshmuminov D, Aminjonov B, Palm RF, Malleo G, Schmockler RK, Abdallah R, et al. FOLFIRINOX or Gemcitabine-based Chemotherapy for Borderline Resectable and Locally Advanced Pancreatic Cancer: A Multi-institutional, Patient-Level, Meta-analysis and Systematic Review. *Ann Surg Oncol.* 2023; 30 (7): 4417–28. [DOI: 10.1245/s10434-023-13353-2] [PMID: 37020094]
18. Conroy T, Castan F, Lopez A, Turpin A, Ben Abdelghani M, Wei AC, et al. Canadian Cancer Trials Group and the Unicancer-GI-PRODIGE Group. Five-Year Outcomes of FOLFIRINOX vs Gemcitabine as Adjuvant Therapy for Pancreatic Cancer: A Randomized Clinical Trial. *JAMA Oncol.* 2022; 8 (11): 1571–8. [DOI: 10.1001/jamaoncol.2022.3829] [PMID: 36048453]
19. Ma SJ, Serra LM, Bartl AJ, Han HR, Fekrmandi F, Iovoli AJ. Adjuvant chemotherapy versus observation following neoadjuvant therapy and surgery for resectable stage I-II pancreatic cancer. *J Radiother Pract.* 2022; 21 (3): 383–92. [DOI: 10.1017/s1460396921000194] [PMID: 36016861]
20. Chan KKW, Guo H, Cheng S, Beca JM, Redmond-Misner R, Isaranuwatthai W, et al. Real-world outcomes of FOLFIRINOX vs gemcitabine and nab-paclitaxel in advanced pancreatic cancer: A population-based propensity score-weighted analysis. *Cancer Med.* 2020; 9 (1): 160–9. [DOI: 10.1002/cam4.2705] [PMID: 31724340]

21. Levine JM, Rompen IF, Franco JC, Swett B, Krysch MC, Habib JR. The impact of metastatic sites on survival rates and predictors of extended survival in patients with metastatic pancreatic cancer. *Pancreatology*. 2024; S1424-3903(24)00659-8. [DOI: 10.1016/j.pan.2024.06.004] [PMID: 38969544]
22. Brown TJ, Reiss KA, O'Hara MH. Advancements in Systemic Therapy for Pancreatic Cancer. *Am Soc Clin Oncol Educ Book*. 2023; 43: e397082. [DOI: 10.1200/EDBK_397082] [PMID: 37192430]
23. Katayama ES, Hue JJ, Bajor DL, Ocuin LM, Ammori JB, Hardacre JM. A comprehensive analysis of clinical trials in pancreatic cancer: what is coming down the pike? *Oncotarget*. 2020; 11 (38): 3489–501. [DOI: 10.18632/oncotarget.27727] [PMID: 33014285]
24. Golan T, Hammel P, Reni M, Van Cutsem E, Macarulla T, Hall MJ, et al. Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer. *N Engl J Med*. 2019 Jul; 381 (4): 317–27. [DOI: 10.1056/NEJMoa1903387] [PMID: 31157963]
25. Maio M, Ascierto PA, Manzyuk L, Motola-Kuba D, Penel N, Cassier PA, et al. Pembrolizumab in microsatellite instability high or mismatch repair deficient cancers: updated analysis from the phase II KEYNOTE-158 study. *Ann Oncol*. 2022;33 (9): 929–38. [DOI: 10.1016/j.annonc.2022.05.519] [PMID: 35680043]
26. Mileusnić D, Marošević G, Durbaba M. *Radijaciona onkologija*. Medicinski fakultet Univerziteta u Banja Luci; 2020. ISBN 978-99976-26-39-4.
27. Versteijne E, van Dam JL, Suker M, Janssen QP, Groothuis K, Akkermans-Vogelaar JM, et al. Dutch Pancreatic Cancer Group. Neoadjuvant Chemoradiotherapy Versus Upfront Surgery for Resectable and Borderline Resectable Pancreatic Cancer: Long-Term Results of the Dutch Randomized PREOPANC Trial. *J Clin Oncol*. 2022; 40 (11): 1220–30. [DOI: 10.1200/JCO.21.02233] [PMID: 35084987]
28. Falco M, Masojć B, Sulikowski T. Radiotherapy in Pancreatic Cancer: To Whom, When, and How? *Cancers (Basel)*. 2023; 15 (13): 3382. [DOI: 10.3390/cancers15133382] [PMID: 37444492]
29. Fietkau R, Ghadimi M, Grützmann R, Wittel U, Jacobasch L, Uhl W, et al. Randomized phase III trial of induction chemotherapy followed by chemoradiotherapy or chemotherapy alone for nonresectable locally advanced pancreatic cancer: First results of the CONKO-007 trial. *Journal of Clinical Oncology*. 2022; 40 (16): 4008-4008. [DOI: 10.1200/JCO.2022.40.16_suppl.4008]
30. Pappalardo A, Giunta EF, Tirino G, Pompella L, Federico P, Daniele B. Adjuvant Treatment in Pancreatic Cancer: Shaping the Future of the Curative Setting. *Front Oncol*. 2021; 11: 695627. [DOI: 10.3389/fonc.2021.695627] [PMID: 34485130]