

ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

Surveillance of influenza in the post-pandemic period in Vojvodina, Serbia, October 2010 – May 2015

Mioljub Ristić^{1,5}, Vesna D. Stojanović^{2,5}, Vesna Milošević^{1,5}, Jelena Radovanov¹, Tihomir Dugandžija^{3,5}, Artur Bjelica^{4,5}, Vladimir Petrović^{1,5}

¹Institute of Public Health of Vojvodina, Novi Sad, Serbia;

²Institute for Child and Youth Health Care of Vojvodina, Novi Sad, Serbia;

³Oncology Institute of Vojvodina, Sremska Kamenica, Serbia;

⁴Clinical Center of Vojvodina, Clinic for Gynecology and Obstetrics, Novi Sad, Serbia;

⁵University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia



SUMMARY

Introduction/Objective In August 2010, World Health Organization declared the beginning of the post-pandemic phase of influenza surveillance.

The aim of this study was to evaluate the epidemiological and virological characteristics of influenza and correlation between the influenza occurrence and weather conditions.

Methods We used surveillance reports of influenza and laboratory data from October 2010 to May 2015. Data for the analysis were collected through sentinel surveillance of influenza-like illness (ILI), severe acute respiratory illness (SARI), acute respiratory distress syndrome, and by virological surveillance. The nasal and throat swabs from all influenza cases were performed by the PCR laboratory method.

Results During the observed period, the highest rates of ILI were registered during the 2010/11 and 2012/13 seasons, with influenza A (H1N1)pdm09 and influenza B being predominant, respectively. The highest weekly age-specific rates of ILI were registered in school-age children (ages 5–14). Out of 1,466 samples collected, 720 (49.1%) were laboratory confirmed as influenza, and influenza A virus was more frequently detected than influenza B. Among confirmed cases of influenza, participation of patients with SARI or ILI was nearly equal (46% vs. 44.1%). There was a weak correlation observed between the decrease in temperature and rainfall and the increase in influenza detection ($p = -0.04214$ vs. $p = -0.01545$, respectively, $p > 0.05$).

Conclusion There is a need for continuous surveillance in order to predict seasonal trends and prepare for a timely response to influenza outbreak.

Keywords: influenza virus; epidemiology; virology; sentinel surveillance

INTRODUCTION

Increased influenza's viral activity causes a high incidence, exceeding incidence of other infectious diseases [1]. The results of epidemiological surveillance of influenza are important as they provide the information which serve as the early warnings of epidemics, and give us an insight into the type and/or subtype of influenza virus, so we can prepare the adequate influenza vaccines and antiviral drugs. They are also significant for implementing appropriate measures for reducing the number of deaths caused by influenza, hospitalization rates due to complications, and the costs of the treatment of influenza [1, 2, 3].

The main factors which determine the seasonality of influenza are still unclear, especially in the tropical areas of the world. Probably, the combined action of environmental factors (humidity, temperature), factors related to immunity of the population, and demographic factors (population density, population flows, school calendar, and traveling or migration) interfere with viral circulation in different parts of the world [4, 5].

Surveillance of influenza at the European level has been conducted since 1996 [6]. In order to improve the monitoring of the epidemiological situation, to ensure efficient response and to reduce the negative consequences on the health of the population, sentinel surveillance of influenza like illness (ILI) and acute respiratory infection (ARI) was introduced in the Autonomous Province of Vojvodina (the northern region of Serbia) in the 2004/05 season, as a pilot study, according to the model of surveillance conducted in Slovenia [7]. In the preparation for an influenza pandemic, sentinel surveillance in Vojvodina became the standard model and also a part of a newly established national influenza surveillance, throughout the Republic of Serbia, since 2009. Due to the quality of the results of the surveillance of influenza in Vojvodina, special Public Health Program was introduced for surveillance of ILI/ARI, severe acute respiratory illness (SARI), and acute respiratory distress syndrome (ARDS) [8]. Similar to other countries in Europe, Vojvodina, as part of Serbia, has a reference laboratory for influenza. In this way, virological surveillance as an integral part of

Received • Примљено:

September 1, 2016

Revised • Ревизија:

October 27, 2016

Accepted • Прихваћено:

November 7, 2016

Online first: March 14, 2017

Correspondence to:

Mioljub RISTIĆ
Institute of Public Health of
Vojvodina
Futoška 121, 21000 Novi Sad
Serbia
mioljub.ristic@mf.uns.ac.rs

Table 1. Number of observed inhabitants during the sentinel surveillance of influenza-like illness by five seasons in the Autonomous Province of Vojvodina

Age groups Seasons	0–4	5–14	15–29	15–64	30–64	≥ 65	Total	Number of included sentinel physicians
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
2010/11	9,281 (10.5)	15,563 (8.2)	-	47,301 (3.5)	-	27,413 (8.7)	99,558 (5.2)	93
2011/12	9,843 (11.1)	17,857 (9.5)	-	60,001 (4.5)	-	25,565 (8.1)	113,266 (5.9)	94
2012/13	10,388 (11.7)	19,837 (10.5)	23,000 (6.3)	-	55,418 (5.7)	27,651 (8.7)	136,294 (7.1)	89
2013/14	23,664 (26.7)	61,157 (32.4)	56,303 (15.5)	-	102,271 (10.5)	48,352 (15.3)	304,719 (15.8)	135
2014/15	17,064 (19.2)	33,298 (17.6)	34,609 (9.5)	-	66,152 (6.8)	33,420 (10.6)	184,543 (9.6)	135

n – surveillance population by age groups and seasons;

% – percentage of population in certain age group according to the census

epidemiological surveillance provides useful information about circulation of an actual influenza virus [9].

In August 2010, the World Health Organization (WHO) declared the post-pandemic period [10]. However, influenza A (H1N1) pdm09, A (H3N2), and B viruses have continued to circulate in the population after the pandemic period [11].

The main goal of this study was to observe seasonal influenza activity along with influenza occurrence and weather conditions in Vojvodina, based on data collected between October 2010 and May 2015, in the post-pandemic period (five surveillance seasons).

METHODS

Data for this observational study were obtained from the sentinel surveillance of ILI, surveillance of all hospitalized patients with SARI and/or ARDS, and virological surveillance of influenza in Vojvodina through the whole season (from calendar week 40 of each year to calendar week 20 of the next year). All information about patients was anonymized and de-identified. Data on temperature and average rainfall were provided by the Republic Hydrometeorological Service of Serbia.

Sentinel surveillance of ILI

During the seasons of surveillance of ILI and ARI, from 2010 to 2015, sentinel surveillance was conducted in all seven districts of Vojvodina. In the surveillance of ILI and ARI, only outpatients in Health Centers of Vojvodina were covered. In accordance with the WHO and national recommendations, the surveillance of influenza was conducted within four age groups (0–4, 5–14, 15–64, and ≥ 65 years) during the 2010/11 and 2011/12 seasons. Since the 2012/13 influenza season, the surveillance has included five age groups (0–4, 5–14, 15–29, 30–64, and ≥ 65 years). Between the 2010 and 2013 seasons, sentinel surveillance has been implemented in 19 municipalities with more than 30,000 inhabitants, encompassing between 5.2% and 7.1% of the total population. During the last two seasons (2013–2015), sentinel surveillance of influenza was ex-

tended to municipalities with less than 30,000 inhabitants and covered all of 45 municipalities of the Province (and included 135 sentinel physicians). Although the number of sentinel physicians during the two last seasons was the same, because we included different sentinel physicians, the observed populations were not equal (Table 1).

The surveillance system used the network of sentinel physicians (general practitioners and pediatricians) who reported the number of new cases of ILI in their reference populations weekly, and collected respiratory samples for virological evaluations. In addition, sentinel physicians electronically entered the data of new cases of ILI and ARI by week, and regularly sent samples for virological confirmation to the WHO National Influenza Center, the Center of Virology of the Institute of Public Health of Vojvodina in Novi Sad. Each week, global population was adjusted with ILI and ARI rate for Vojvodina, which allows to follow the indicators of influenza activity [7]. The influenza surveillance data were collected for the whole season (from calendar week 40 of each year to calendar week 20 of the following year) [12].

SARI and ARDS surveillance

Since the 2010/11 influenza season, in accordance with the WHO recommendations, SARI surveillance has been carried out during the winter season only, from week 40 to week 20 [13].

As described in detail previously, hospital coordinators for SARI and ARDS surveillance from all 15 acute care hospitals in Vojvodina sent daily reports on every hospitalized SARI and/or ARDS case to the district coordinators of influenza in the local departments of Public Health [7]. The study included patients from intensive care units and high dependency units (severe form of infections), general/internal medicine, pediatric medicine, infectious disease wards and respiratory disease wards where patients of all ages were registered [8].

Case definitions

This study included all patients with ILI and any flu-associated infections who required hospitalization in accordance

with the SARI and ARDS definitions. The WHO criteria for ILI and SARI were used for screening in the primary care/outpatient settings and inpatient hospital settings: ILI cases were defined as those with a sudden onset of fever ($> 38^{\circ}\text{C}$) and cough/sore throat within seven days of the onset, while patients with an acute respiratory illness with the onset of symptoms during the previous seven days requiring overnight hospitalization which included history of fever or measured fever of 38°C , and cough, and shortness of breath or difficulty in breathing were termed as SARI [14].

ARDS cases were defined as acute onset of bilateral infiltrates on the chest radiograph; arterial oxygen tension (PaO_2)/fraction of inspired oxygen (FiO_2) ratio $< 27\text{kPa}$, and absence of a cardiac failure or left atrial hypertension (assessed clinically, echocardiographically, or with invasive monitoring) and required invasive ventilation [15, 16, 17].

Virological surveillance

The reference laboratory for virological surveillance of influenza in Vojvodina is the WHO National Influenza Center at the Institute of Public Health of Vojvodina, Novi Sad [18].

Nasal and throat swabs from patients with suspected influenza were placed in the same vial with transport medium and kept at -20°C . The transport of the samples to the laboratory was organized on a daily basis by local departments of public health. Viral RNA was extracted using the commercially available QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The reverse transcription and amplification were performed using the AgPath-IDTM One-Step RT-PCR Reagents (Applied Biosystems, Foster City, CA, USA) on a 7500 Real-Time (Applied Biosystems) thermocycler. Influenza A and B virus detection (without further determination of B/Yamagata-like and B/Victoria-like viruses) was done by singleplex real-time RT-PCR assays. The testing was done according to the protocol developed by the Centers for Disease Control and Prevention, enclosed with the reagents. The results were analyzed using the Applied Biosystems 7500 Software version 2.0.6, and the interpretation of the data was done according to the WHO guidelines [19]. Immediately after the testing was finished, the results of the laboratory tested samples were sent to the Institute of Public Health of Serbia in Belgrade, local departments of public health, the sentinel/hospital physician, and to the patients [7].

Statistical analysis

Similar to the previously used methodology, the population under surveillance was used as a denominator for calculations of the weekly incidence of ILI and the numerator was the number of clinical cases of ILI in the total population and in the age groups [7]. Also, the weekly age-specific incidences of ILI for monitored age groups were measured per 100,000 inhabitants.

The epidemic threshold of incidence of $246.3/100,000$ was determined in the previous five pre-pandemic sentinel

seasons on the basis of the weekly incidence rate of ILI value [7].

A correlation analysis was performed using the Spearman correlation coefficient between the detected monthly influenza cases and the average monthly temperature and rainfall using the data from the Republic Hydrometeorological Service of Serbia, for the capital of the Autonomous Province of Vojvodina, city of Novi Sad [20]. Because the Meteorological annual data from the Republic Hydrometeorological Service of Serbia contains the average monthly values for only six different meteorological stations (regions) in Serbia (Belgrade, Novi Sad, Vranje, Zlatibor, Loznica, and Niš), we used values of average monthly temperature and rainfall only for the city of Novi Sad (Vojvodina), which represents weather conditions for this region of Serbia. Average temperature and rainfall were measured in degrees Celsius ($^{\circ}\text{C}$) and millimeters (mm), respectively.

Two tailed p-values less than 0.05 were considered statistically significant.

RESULTS

Surveillance of ILI

During the study period (2010–2015), the highest weekly incidence rates were between $474.5/100,000$ (2010/11 season, with influenza A (H1N1)pdm09 being predominant) and $712.3/100,000$ (2012/13 season, with influenza B being predominant). The lowest weekly incidence rate was registered during the 2013/14 season, predominated by influenza A (H3N2). In the 2010/11 influenza season, with the predominance of influenza A (H1N1)pdm09 (ILI incidence above the epidemic threshold of 246.3 cases per 100,000 population) the duration of the epidemic period was eight weeks. Only during the 2013/14 season, values of the weekly ILI incidence were below the epidemic threshold (Figure 1).

Figure 2 shows the trend of ILI weekly incidence rate by age group during the five post-pandemic seasons. During the observed period, it was evident that the highest weekly age-specific incidence of ILI was registered in school-age children (5–14 years old), with the highest incidence rate




Figure 1. Influenza-like illness (ILI) in the 2010–15 seasons according to the influenza sentinel surveillance system in the Autonomous Province of Vojvodina




Figure 2. Age-specific incidence of ILI per 100,000 population in the Autonomous Province of Vojvodina during five influenza seasons; black vertical lines indicate the separation between different influenza seasons (2010–15)




Figure 3. Incidence rate of ILI consultations per week by case status for the sentinel surveillance network, in the Autonomous Province of Vojvodina, during 5 influenza seasons; black vertical lines indicate the separation between different influenza seasons (2010–15)

Table 2. Distribution of influenza infection by case definitions and by type/subtype in the Autonomous Province of Vojvodina, 2010–15 seasons

Variable	2010/11		2011/12		2012/13		2013/14		2014/15		Total		
	n	%	n	%	n	%	n	%	n	%	n	%	
Clinical form	ILI	29	24.6	49	92.5	72	36.2	63	53.8	105	45.1	318	44.1
	SARI	66	55.9	4	7.5	110	55.3	47	40.2	104	44.6	331	46
	ARDS	23	19.5	0	-	17	8.5	7	6	24	10.3	71	9.9
Influenza virus type/subtype	A (H1N1)pdm09	101	85.6	0	-	68	34.2	33	28.2	86	36.9	288	40
	A (H3N2)	6	5.1	52	98.1	28	14.0	77	65.8	90	38.6	253	35.1
	A (non-subtype)	3	2.5	0	-	2	1.0	7	6	9	3.9	21	3
	B	8	6.8	1	1.9	101	50.8	0	-	48	20.6	158	21.9
Total	118	100	53	100	199	100	117	100	233	100	720	100	

ILI – influenza-like illness, SARI – severe acute respiratory infection; ARDS – acute respiratory distress syndrome

of 2,315.2/100,000 in the 9/2013 week. In all five seasons, the lowest weekly age-specific incidence of ILI was registered in the oldest population.

In all five influenza seasons, the highest value of weekly ILI incidence rate was accompanied by the highest number of suspected and confirmed influenza cases (Figure 3).

Influenza by case definitions and prevalence distribution of influenza viruses

In the observed period, 1,466 specimens from patients with ILI, SARI, or ARDS, were tested, and 720 samples were identified as influenza A or B positive (49.1%). There were a total of 562 influenza A and 158 influenza B virus infections confirmed during the five influenza seasons. During all five seasons, influenza A (H1N1)pdm09 was predominant (40%, 288/720). Participation of patients with SARI or ILI was nearly equal (46%, 331/720 vs. 44.1%, 318/720, respectively). Patients with ARDS were not registered only during the 2011/12 influenza season (Table 2).

Weather conditions and prevalence of influenza

The dependence of influenza activities on weather conditions in the Province (meteorological station for the city of Novi Sad) during the investigation period is shown in Figure 4. During three influenza seasons (2011/12, 2012/13, and 2014/15), the highest number of confirmed

influenza infections was registered during March. The average monthly temperature (October–April, 2010–15) was between -4.9°C and 13.6°C , and average values of rainfall in the same period were 1.3–82.8 mm. These data show that the average monthly temperature and average rainfall do not have any significant correlation with the number of influenza-positive cases ($\rho = -0.04214$ vs. $\rho = -0.01545$; $p > 0.05$, respectively).

DISCUSSION

We presented the first description of the epidemiological and virological characteristics of influenza and the correlation between the influenza occurrence and weather conditions after the pandemic 2009/10 season in Vojvodina. As can be concluded from the obtained data, all influenza viruses were detected across Vojvodina and all age groups were affected during the post-pandemic seasons.

During the 2010/11 season in Western Europe, transmission peaked during late January and early February, and two to three weeks later in Eastern Europe, with influenza A (H1N1) virus as predominant in 2009. Opposite Europe, where influenza A (H3N2) was rare, it was predominant in North America. During this season, in several countries, a higher number of severe cases of influenza were registered compared to the previous season, but reasons for this were not clear [21]. As it was expected, influenza A (H1N1)pdm09 virus predominated among the vi-




Figure 4. Correlation of the number of influenza cases in the Autonomous Province of Vojvodina in the investigated period with: (a) the average temperature (°C); (b) the average rainfall (mm); black vertical lines indicate the separation between different influenza seasons (2010–15)

ruses detected during the first post-pandemic season, but the results of the phylogenetic analysis of HA sequences of A(H1N1)pdm09 viruses circulating in Lombardy showed that viruses isolated during the 2010/11 season segregated into a different genetic group with respect to those identified during the 2009 pandemic [22].

In Vojvodina, during the 2010/11 season, similar to the previous season, influenza A (H1N1)pdm09 was predominant (85.6%, 101/118), with peaked transmission during late January and throughout all of February. The highest incidence rate of ILI was registered in the 5–14 years age group. The epidemic wave (incidence rate of ILI above 246.3/100,000 population) lasted for eight weeks, and as much as 75.4% (89/118) of all confirmed cases of influenza were hospitalized for SARI or ARDS.

Similar to the results of the WHO review, in Vojvodina, during the 2011/12 season, influenza transmission peaked during late March [23]. Influenza A (H3N2) was predominant 98.1% (52/53), and the highest incidence rate of ILI was registered among the youngest patients and in the school-age children (0–4 and 5–14 years old). The epidemic wave lasted for only three weeks.

In North America, the 2012/13 influenza season started earlier than in other parts of the Northern Hemisphere. In Europe, the influenza season peaked two weeks later than in North America, and it was also unusually long, associated with a late rise in influenza B cases in many countries. Peaking of influenza activity was registered around week 5/2013 and lasted until week 16/2013. In North America and Europe, later in the season, influenza B type was more commonly detected than either of the influenza A subtypes [24, 25].

In Vojvodina, the highest incidence rate of ILI was registered in the 9/2013 week (712.3/100,000 population), and the epidemic wave lasted for six weeks (weeks 6–11). Among confirmed cases of influenza, the predominant type was influenza B (50.8%, 101/199).

In the 2013/14 season in Europe, influenza activity started later than usual, with a very small increase during the final weeks of December, followed by more marked increase throughout January. Overall activity continued to increase through February. Influenza activity during the 2013/14 season was less intense, with fewer positive sam-

ples detected than in the previous post-pandemic seasons. Overall, influenza A (H1N1)pdm09 was predominant in most of northern European countries and A (H3N2) was predominant in most of eastern European countries. Unlike North America, where influenza B was detected in the late season, the detection of influenza B in Europe remained low during the entire season [26].

In our territory, during the 2013/14 influenza season, the first influenza case was registered in the 2/2014 week, and the incidence rate of ILI was below the baseline for the entire season. Similarly to the 2011/12 influenza season, the highest incidence rate of ILI was registered among the youngest and in the school-age children. We think that the reasons for the highest incidence rates of ILI among the mentioned age groups perhaps lie in the fact that young people are tested more often [7]. Furthermore, as is well known, children may shed viruses more profusely and for longer periods of time than adults [19].

Similarly to the results for Eastern Europe, the predominant subtype of influenza in Vojvodina was A (H3N2), with 65.8% (77/117) of all confirmed cases [26]. Unlike the previous season, when influenza B was the predominant type, it was not detected among the tested samples during the 2013/14 season.

Considering European countries, the timing of influenza detections during the 2014/15 season was similar to the previous years. The influenza activity started increasing in the last few weeks of 2014, similar to the situations in the 2013/14 and 2011/12 seasons. The peak of influenza activity in 2015 varied between countries, but it was most often between weeks 6 and 9. However, the percentage of positive influenza cases was still above the threshold in many countries in April. During the 2014/15 season, influenza A (H3N2) virus was predominant in most regions, and an increased proportion of influenza B virus was registered after the activity peak. During this influenza season, most of circulating influenza A (H3N2) viruses were different from the virus used in vaccines in the northern hemisphere [27, 28].

The data obtained from this study showed that during the 2014/15 season, epidemic influenza activity started in the 6/2015 week and lasted to the 11/2015 week. Like in previous seasons, the lowest weekly age-specific

incidence of ILI was registered in the oldest population. This phenomenon can be interpreted by the fact that older patients are more likely to be registered in the hospital institutions than at the primary care level (sentinel surveillance of influenza). During the last influenza season, the predominant virus was A (H3N2), with 38.6% (90/233), whereas the registered proportion of influenza B was 20.6% (48/233) of all confirmed cases. The last cases of influenza were registered on April 30, 2015.

In terms of seasonal pattern, in the Northern Hemisphere, influenza viruses are more frequently detected in autumn, winter, and spring, with the peak of influenza activity occurring typically between December and March and lasting for six to eight weeks. In temperate regions of the Southern Hemisphere, influenza activity typically peaks between May and September. In addition, influenza viruses can be isolated sporadically throughout the year [19].

In countries with tropical climate, influenza activity has been reported to peak in the rainy season, between June and August [29].

The climate in our country can be described as moderate continental, with intermediate temperatures and relatively low humidity. June is the rainiest, with the average of 12–13% of total annual precipitation sum. February and October have the least of precipitation [20]. During the study period (from the beginning of October to the end of April, 2010–2015) there was no significant correlation between the number of influenza detections by months and average monthly temperature and rainfall.

Although almost 55% of all confirmed cases of influenza were recorded in the city of Novi Sad, absence of data

on monthly average temperature and rainfall in other parts of the Province and their comparison with influenza cases may be considered to be a limitation of this study.

CONCLUSION

The findings of the presented sentinel ILI and SARI surveillance, along with the virological surveillance, which are comparable with the results of surveillance of influenza in other countries across Europe, offer the possibility to determine the epidemiology of influenza during the post-pandemic seasons and predict future epidemic occurrences of influenza in Vojvodina. Likewise, the increasing number of samples among some age groups (0–4- and 5–14-year-olds) with the highest value of weekly ILI incidence rate would lead to a more precise view on the distribution of influenza in Vojvodina in the younger population.

ACKNOWLEDGMENTS

The authors are grateful to all physicians who participated in the influenza surveillance system in Vojvodina during the 2010–2015 seasons.

NOTE

This investigation was carried out as part of routine activities without additional funding sources.

REFERENCES

- Aguilera JF, Paget WJ, Mosnier A, Heijnen ML, Uphoff H, van der Velden J, et al. Heterogeneous case definitions used for the surveillance of influenza in Europe. *Eur J Epidemiol*. 2003; 18(8):751–4.
- World Health Organization. The role of National Influenza Centres (NICs) during inter-pandemic, pandemic alert and pandemic periods. Interim Document. 2007:1–13. Available from: http://www.who.int/csr/disease/avian_influenza/guidelines/RoleNICsMayf.pdf
- World Health Organization, Department of Communicable Disease Surveillance and Response. WHO global influenza preparedness plan. The role of WHO and recommendations for national measures before and during pandemics. WHO/CDS/CSR/GIP/2005.5, Global Influenza Programme, World Health Organization, Geneva, Switzerland, 2005:1–50.
- Tamerius J, Nelson MI, Zhou SZ, Viboud C, Miller MA, Alonso WJ. Global influenza seasonality: reconciling patterns across temperate and tropical regions. *Environ Health Perspect*. 2011; 119(4):439–45.
- Freitas FT. Sentinel surveillance of influenza and other respiratory viruses, Brazil, 2000–2010. *Braz J Infect Dis*. 2013; 17(1):62–8.
- European Influenza Surveillance Scheme. Annual Report 2005–2006 influenza season. Utrecht, the Netherlands, NIVEL, 2007:1–38.
- Petrović V, Seguljev Z, Cosić G, Ristić M, Nedeljković J, Dragnić N, et al. Overview of the winter wave of 2009 pandemic influenza A(H1N1)v in Vojvodina, Serbia. *Croat Med J*. 2011; 52(2):141–50.
- Institute of Public Health of Vojvodina. Communicable diseases in Vojvodina, 2015. Annual report. Novi Sad; 2016.
- Losos JZ. Routine and sentinel surveillance methods. *East Mediterr Health J*. 1996; 2:46–50.
- Eurosurveillance editorial team. World Health Organization declares post-pandemic phase. *Euro Surveill*. 2010; 15(32).
- Guan WD, Gong XY, Mok CK, Chen TT, Wu SG, Pan SH, et al. Surveillance for seasonal influenza virus prevalence in hospitalized children with lower respiratory tract infection in Guangzhou, China during the post-pandemic era. *PLoS One*. 2015; 10:e0120983.
- Ministry of Health of the Republic of Serbia. New guidelines for A(H1N1)v influenza surveillance in the Republic of Serbia.
- World Health Organization. Overview of sentinel systems for hospitalized severe acute respiratory infections (SARI) represented in the weekly EuroFlu surveillance bulletin (as of 10 February 2013, p. 20). [accessed 2013 July 15] Available from: http://www.euro.who.int/__data/assets/pdf_file/0005/186863/Overview-of-SARISurveillance-Systems-final.pdf.
- World Health Organization (2009). Human infection with pandemic (H1N1) 2009 virus: updated interim WHO guidance on global surveillance. Available from: http://www.who.int/csr/disease/swineflu/guidance/surveillance/WHO_case_definition_swine_flu_2009_04_29.pdf?ua=1 [accessed 2014 July 20].
- Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, et al. The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med*. 1994; 149:818–24.
- Institut za javno zdravlje Srbije. Stručno-metodološko uputstvo za kontrolu unošenja i sprečavanja širenja novog soja virusa gripa A(H1N1) – revizija 1.
- Irish Critical Care Trials Group. Acute lung injury and the acute respiratory distress syndrome in Ireland: a prospective audit of epidemiology and management. *Crit Care*. 2008; 12:R30.
- World Health Organization. National Influenza Centres. Institute of Public Health; Serbia – Novi Sad. [accessed 2015 September 12] Available from: http://www.who.int/influenza/gisrs_laboratory/national_influenza_centres/list/en/index3.html.
- World Health Organization. Global Influenza Surveillance Network - WHO GISN. Manual for the laboratory diagnosis and virological surveillance of influenza. Geneva, Switzerland: WHO Press; 2011.

20. Republic Hydrometeorological Service of Serbia (RHMS). [accessed 2015 July 15]. Available from: <http://www.hidmet.gov.rs/eng/meteorologija/klimatologija.php>.
21. World Health Organization. Review of the 2010–2011 winter influenza season, northern hemisphere. *Wkly Epidemiol Rec.* 2011; 86(22):222–7.
22. Pariani E, Amendola A, Ranghairo A, Anselmi G, Zanetti A. Surveillance of influenza viruses in the post-pandemic era (2010–2012) in Northern Italy. *Hum Vaccin Immunother.* 2013; 9(3):657–66.
23. World Health Organization. Review of the 2011–2012 winter influenza season, northern hemisphere. *Wkly Epidemiol Rec.* 2012; 87(24):233–40.
24. World Health Organization. Review of the 2012–2013 winter influenza season, northern hemisphere. *Wkly Epidemiol Rec.* 2013; 88(22):225–32.
25. World Health Organization. Summary of the 2012–2013 influenza season in the WHO European Region. [accessed 2015 July 9]. Available from: http://www.euro.who.int/__data/assets/pdf_file/0011/195824/Summary-of-the-20122013-influenza-season-in-the-WHO-European-Region-final.pdf
26. World Health Organization. Review of the 2013–2014 winter influenza season, northern hemisphere. *Wkly Epidemiol Rec.* 2014; 89(23):245–56.
27. World Health Organization. Review of the 2014–2015 influenza season in the northern hemisphere. *Wkly Epidemiol Rec.* 2015; 90(23):281–96.
28. World Health Organization. Regional Office for Europe influenza season summaries. [accessed 2015 April 10]. Available from: <http://www.euro.who.int/en/health-topics/communicable-diseases/bulletins-and-seasonal-summaries/influenza-season-summaries2>
29. Prachayangprecha S, Makkoch J, Suwannakarn K, Vichaiwattana P, Korkong S, Theamboonlers A, et al. Epidemiology of seasonal influenza in Bangkok between 2009 and 2012. *J Infect Dev Ctries.* 2013; 7(10):734–40.

Надзор над инфлуенцом током постпандемијског периода у Војводини, од октобра 2010. до маја 2015.

Миољуб Ристић^{1,5}, Весна Д. Стојановић^{2,5}, Весна Милошевић^{1,5}, Јелена Радованов¹, Тихомир Дуганџија^{3,5}, Артур Бјелица^{4,5}, Владимир Петровић^{1,5}

¹Институт за јавно здравље Војводине, Нови Сад, Србија;

²Институт за здравствену заштиту деце и омладине Војводине, Нови Сад, Србија;

³Институт за онкологију Војводине, Сремска Каменица, Србија;

⁴Клинички центар Војводине, Клиника за гинекологију и акушерство, Нови Сад, Србија;

⁵Универзитет у Новом Саду, Медицински факултет, Нови Сад, Србија

САЖЕТАК

Увод/Циљ У августу 2010. године Светска здравствена организација је прогласила почетак постпандемијске фазе у надзору над gripом.

Циљ рада био је да се процене епидемиолошке и вирусолошке карактеристике gripа и однос између климатских услова и преваленције gripа у постпандемијском периоду.

Методe Коришћени су подаци из надзора над gripом и резултати лабораторијских испитивања у периоду од октобра 2010. до маја 2015. године. Подаци су прикупљени кроз сентинелни надзор за обољења слична gripу (ОСГ), тешку акутну респираторну болест (ТАРБ), акутни респираторни дистрес синдром (АРДС) и вирусолошким надзором. Сви случајеви gripа потврђени су из назофарингеалног бриса лабораторијском методом *PCR*.

Резултати У посматраном периоду највише стопе ОСГ регистроване су током сезоне 2010/11. и 2012/13, када су

доминирали вирус gripа типа А (*H1N1*)*pdm09*, односно вирус gripа типа Б. Највише вредности узрасно специфичних стопа ОСГ регистроване су у школском узрасту (5–14 година). Од 1.466 тестираних узорака, лабораторијска потврда вируса gripа добијена је код 720 (49,1%), а вирус gripа типа А чешће је детектован од вируса gripа типа Б. Међу потврђеним случајевима gripа учешће оболелих са дијагнозама ТАРБ или ОСГ је било скоро подједнако (46 тј. 44,1%). Није утврђена статистички значајна разлика између смањења температуре и падавина у односу на пораст броја потврђених случајева gripа ($p = -0,04214$ наспрам $p = -0,01545$, $p > 0,05$).

Закључак Наставак надзора над gripом кроз процену сезонских трендова омогућава спремност на одговор у случају појаве епидемије gripа.

Кључне речи: вирус gripа; епидемиологија; вирусологија; сентинелни надзор