



ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

Complications after total thyroidectomy in correlation with histopathological and hormonal findings

Mirjana Gnjačić^{1,2}, Dalibor Vranješ^{1,2}, Vanja Nežić², Branko Jurišić^{1,2}, Irma Ovčina³

¹University of Banja Luka, Faculty of Medicine, Banja Luka, Republic of Srpska, Bosnia and Herzegovina;

²University Clinical Center of Republic of Srpska, ENT Clinic, Banja Luka, Republic of Srpska, Bosnia and Herzegovina;

³University Clinical Center of Republic of Srpska, Clinic for Internal Diseases, Banja Luka, Republic of Srpska, Bosnia and Herzegovina

SUMMARY

Introduction/Objective Total thyroidectomy (TT) is frequently performed in head and neck surgery and is considered a relatively straightforward procedure, yet it is still associated with a number of complications. This study aimed to examine the role and significance of TT in the occurrence of postoperative complications in correlation with histopathological (HP) and endocrinological findings.

Methods A cross-sectional study was conducted involving 90 patients in the University Clinical Center of the Republic of Srpska, Banja Luka, Republic of Srpska, Bosnia and Herzegovina who underwent the thyroid gland surgery 2022–2024. Preoperatively, hormonal analyses, thyroid ultrasound examination, and endovideolaryngoscopy were performed. TT was performed in all patients. Postoperatively, HP diagnostics, control hormonal tests, and endovideolaryngoscopy were performed.

Results Vocal fold paralysis (VFP) was reported by 11 (12.2%) patients. HP analysis showed that goiter was the leading cause in 37 (41.1%) patients, followed by follicular adenoma in 25 (27.8%) and papillary carcinoma in 18 (20%); other causes were present in the remaining 10 (11%). Regarding endocrinological complications, hypocalcemia was present in 67 (74.4%) patients, and elevated parathyroid hormone levels in four (4.4%) patients.

Conclusion VFP and hypocalcemia represented the most common postoperative complications after TT, particularly in patients with HP-diagnosed goiter, follicular adenoma, or papillary thyroid carcinoma. These findings may help identify older patients who are at increased risk of TT-related complications that could significantly impair their quality of life.

Keywords: hypocalcemia; thyroid gland diseases; vocal fold paralysis

INTRODUCTION

Thyroid disorders are among the most common endocrine-system diseases, and thyroidectomy is the therapy of choice for benign lesions, and is a surgical intervention routinely used in cases of suspected malignancy after ultrasound examination and fine-needle aspiration [1]. Other reasons for thyroid surgery include enlargement of the thyroid gland in the form of nodular or colloid goiter that interferes with breathing or swallowing [2]. Even with recent adjustments to total thyroidectomy (TT) meant to reduce the financial strain on hospitals, expenses are still growing by 4.3% annually [3]. Currently, this surgical intervention is very common and is among the safest surgical procedures [4]. Ultrasound is a mandatory diagnostic procedure for thyroid disease due to its reliability, non-invasiveness among diagnostic procedures, and its reliability and non-invasiveness, whereas computed tomography and magnetic resonance imaging are reserved for complex cases [5].

However, thyroid surgery still carries a number of complications, regardless of whether it is performed using the classic open technique

or micro-invasive surgery, and even includes a certain degree of mortality risk [6]. The most common complications include hypoparathyroidism (HPT), damage to the external branch of the superior laryngeal nerve, the recurrent laryngeal nerve, postoperative hemorrhage, laryngeal edema, and tracheospasm with dyspnea [6]. Post-thyroidectomy hypocalcemia occurs due to HPT, which can result from the devascularization or unintentional removal of the parathyroid glands [7]. A frequent and dangerous complication after TT is hypocalcemia [8]. Its incidence is about 1.2%, and it is transient in approximately 40% of cases and permanent in only 3% [7, 8]. Another common complication is damage to the recurrent laryngeal nerve.

The incidence of transient recurrent vocal-fold paralysis (VFP) ranges 5–8%, while the rate of permanent recurrent VFP is 0.3–3% [9]. The increase in the number of TT is associated with more postoperative complications. Surgeon experience, or lack thereof, accounts for 46–56% of cases of unilateral and bilateral VFP, which can ultimately lead to dysphonia, dysphagia, and dyspnea [10]. Postoperative dysphonia can

Received • Примљено:
February 24, 2025

Revised • Ревизија:
April 17, 2025

Accepted • Прихваћено:
April 25, 2025

Online first: April 29, 2025

Correspondence to:

Mirjana GNJATIĆ
ENT Clinic
University Clinical Center of
Republic of Srpska
12 beba bb
78000 Banja Luka
Republic of Srpska
Bosnia and Herzegovina
mgnjatic@gmail.com

have significant implications for patients, so it is crucial to take all necessary measures to minimize this complication [11]. Postoperative hypocalcemia has a variety of causes. Since many of these characteristics may be changed, it is critical to identify them after surgery so that high-risk population groups can be identified and early preventive measures can be put in place [12].

For patients who are experiencing changes in their voice, a multidimensional voice assessment and a postoperative laryngeal examination are advised [13]. After thyroid surgery, 80% of patients have voice disorders, meaning that roughly 10% of patients have recurrent laryngeal nerve damage [10]. Dysphonia after TT can significantly affect a patient's quality of life and interfere with their ability to perform work and daily activities [11]. Preoperative voice education, managing anticipated voice changes during surgery, and thorough voice care after TT are all recommended by the guidelines [13]. This study aimed to examine the role and significance of TT in the occurrence of postoperative complications in correlation with histopathological (HP) and endocrinological findings.

METHODS

A cross-sectional study was conducted involving 90 patients with thyroid diseases who underwent TT at the University Clinical Center of the Republic of Srpska (UCCRS) in Banja Luka 2022–2024. Inclusion criteria were that the patients had a thyroid disease not surgically treated until arrival at the center. Patients under 18 years of age and those without a history of fine-needle aspiration were excluded.

The patients underwent an initial examination. Demographic characteristics and medical history were recorded at the Clinic for Endocrine Diseases. Indications for surgery were determined by endocrinologists who, based on anamnestic data, clinical examination, ultrasonography of the thyroid gland, laboratory findings in the form of hormonal status and basic laboratory tests, diagnosed one of the thyroid diseases. Following TT, which involved removing all thyroid tissue, calcium, parathyroid hormone (PTH) and FT4 levels were measured at the Clinic for Endocrinology. After 48 h, the patients were examined in the phoniatrics office to assess the condition of the vocal folds, and endovideolaryngoscopy was performed to diagnose whether there was proper mobility of both vocal folds or unilateral or bilateral damage to the *n. laryngeus recurrens*.

Data analysis was conducted using SPSS v22.0 (IBM, USA). Descriptive statistics are presented as the mean, median and standard deviation. Quantitative values are expressed as absolute numbers (n) and percentages (%). For analysis of categorical variables, we used the χ^2 test and Fisher's exact test. Depending on the normality of the data distribution, we used the Student's t-test for independent samples or the Mann-Whitney test. For dependent variables, we used the t-test for dependent samples or the Wilcoxon test. For numerical variables, either Spearman's or Pearson's correlation was used. Results were considered significant if $p < 0.05$.

Ethics: The study was approved by the Ethics Committee of UCCRS (approval no. 01-19-439-2/24) and complied with the revised ethical guidelines of the Declaration of Helsinki.

RESULTS

This cross-sectional study included 90 patients of both sexes. Demographic data, indications for surgery, symptoms before and after surgery, thyroid-stimulating hormone (TSH) before and after surgery, postoperative serum calcium level, PTH and occurrence of unilateral or bilateral vocal-fold paralysis were collected. Of the total number of patients, 81 (90%) were female, with a mean age of 54.18 ± 11.8 years. The most common indication for TT was the isolated presence of goiter in 51 (56.7%), followed by Hashimoto's thyroiditis in 11 (12.2%) (Figure 1).

Five (5.6%) patients had preoperative symptoms, all of whom were women, while 85 (94.4%) patients had no preoperative symptoms. The most common postoperative symptoms were dyspnea on exertion, dizziness, neck pain, dysphagia, and cough. Laryngeal symptoms after TT were present in 13 (14.4%) patients and were reported as: hoarseness; hoarseness and difficulty swallowing; difficulty swallowing and cough (Figure 2). HP findings showed the highest presence of goiter in 37 (41.1%) patients, followed by follicular adenoma in 25 (27.8%) and papillary carcinoma in 18 (20%). VFP occurred in 11 patients (12.2%). The highest incidence of VFP occurred in patients with an HP diagnosis of goiter in 4 (4.4%) patients and papillary carcinoma in 4 (4.4%) patients, but it was not statistically




Figure 1. Frequency of preoperative thyroid disease diagnosis




Figure 2. Presentation of laryngeal symptoms after surgery

significant ($\chi^2 = 3.88$, $df = 3$, $p = 0.27$) (Table 1). In the total sample, 11 (12.2%) had unilateral VFP, while no cases of bilateral VFP were detected.

Table 1. Correlation of pathohistological findings and paralysis

Paresis	HP				p
	Goiter	Papillary carcinoma	Follicular adenoma	Other	
Yes	4 (4.4)	4 (4.4)	1 (1.1)	2 (2.2)	> 0.05
No	33 (36.7)	14 (15.6)	24 (26.7)	8 (8.9)	

$\chi^2 = 3.88$, $df = 3$, $p = 0.27$; HP – histopathological

In the total sample, 12 (13.3%) patients had clinical endocrinological symptoms postoperatively. Among the symptoms reported were calf pain, leg cramps, tingling sensations, and a general feeling of weakness. Sixty-seven (74.4%) patients had hypocalcemia, of which 51 (56.7%) had mild ($1.9\text{--}2.2\text{ mmol L}^{-1}$) and 16 (17.8%) severe hypocalcemia ($< 1.9\text{ mmol L}^{-1}$). Four (4.4%) patients had elevated PTH levels. A statistically significant correlation was found between age and postoperative calcium levels ($r_s = 0.2$, $p = 0.05$). However, no statistically significant associations were observed between age and postoperative TSH ($r_s = -0.09$, $p = 0.37$), age and preoperative TSH ($r_s = 0.004$, $p = 0.97$) or postoperative PTH ($r_s = 0.02$, $p = 0.84$) (Table 2).

Table 2. Correlation of endocrinological complications with laboratory parameters

Laboratory parameters	N	Min.	Max.	Mean	SD
TSH preoperatively	90	0	4.33	1.55	1.14
TSH postoperatively	90	0.01	45	13.84	10.56
PTH postoperatively	90	1	75	32.7	16.28
Hypocalcemia postoperatively	90	1.06	2.56	2.02	0.32

TSH – thyroid-stimulating hormone; PTH – parathyroid hormone

Table 3. Correlation of the occurrence of postoperative symptomatology overall with age and laboratory parameters

	Age	PTH postoperatively	TSH preoperatively	TSH postoperatively
p	0.35	0.005	0.35	0.31

$U = 470$, $p = 0.005$; PTH – parathyroid hormone; TSH – thyroid-stimulating hormone

There is a statistically significant correlation between postoperative PTH levels and the onset of postoperative symptoms overall ($U = 470$, $p = 0.005$). Age does not show a statistically significant correlation with postoperative symptoms ($U = 671.5$, $p = 0.35$), nor do preoperative TSH

($U = 670$, $p = 0.35$), postoperative TSH ($U = 662$, $p = 0.31$) or postoperative pH (Table 3).

Logistic-regression analysis was conducted to evaluate the influence of age, HPT presence, preoperative TSH levels, pH, and sex on the total incidence of postoperative symptoms. The overall model did not show a statistically significant difference when compared with the null model ($\chi^2 = 5.013$, $p = 0.658$), accounting for 8% of the variation in postoperative symptoms (Nagelkerke R^2). There was no statistically significant effect of individual factors on the occurrence of postoperative symptoms: age ($p = 0.45$), hyperthyroidism ($p = 0.68$), preoperative TSH ($p = 0.88$), pH ($p = 0.57$), and sex ($p = 0.20$) (Table 4).

DISCUSSION

The analysis of post-thyroidectomy complications reveals that, while these procedures are typically routine and relatively safe, they do carry some risks. The most common complications, including VFP and HPT, can significantly affect patients' quality of life [14]. TT still has an overall complication rate of up to 54.4% [1]. Our study reveals several key findings regarding laryngeal complications after TT, highlighting their association with HP findings and the patient's hormonal status [9]. The findings also align with earlier research on complications following thyroid surgery, particularly highlighting hypocalcemia and recurrent laryngeal-nerve dysfunction as the most frequently encountered negative outcomes. We demonstrated that the rate of recurrent laryngeal-nerve dysfunction is 12.2%, with symptoms observed in 14.4% of individuals. Our results align with previous studies that have reported comparable complications [15].

Excessive use of alcohol, due to its toxic effect, can be one of the risk factors for thyroid cancer [16]. We found that VFP occurred in 11 (12.2%) patients after TT, with no statistically significant association with sex, age, presence of goiter, or hyperthyroidism. However, goiter and papillary carcinoma were present in the majority of patients with VFP [17]. EVLS offers insights into the anatomical structures and functional changes of the vocal folds, including their appearance and vibratory patterns, without any complications [18]. These results align with research conducted by other authors, which found that a notable proportion of patients experienced VFP, with its occurrence being influenced by

Table 4. Correlation of demographic laboratory and endocrinological and PH factors with postoperative complications

Variable		Coef. (B)	SE	Odds ratio (Exp B)	95% confidence interval		p
					Lower	Upper	
Age		0.016	0.022	0.984	0.943	1.026	0.45
Hyperthyroidism	No			1			
	Yes	0.476	1.159	1.610	0.166	15.592	0.68
TSH before		0.033	0.222	1.034	0.669	1.598	0.88
HP							0.57
1		-0.952	0.794	0.389	0.081	1.828	0.23
2		-0.210	0.846	0.476	0.154	4.257	0.80

χ^2 -test = 5.013, $p = 0.658$, TSH – thyroid-stimulating hormone; HP – histopathological

the surgeon's level of expertise and the scope of the operation [19]. In this regard, the importance of standardizing recurrent laryngeal-nerve identification activities was emphasized to prevent nerve damage during surgery [20].

Regardless of changes in laryngeal mobility, individuals may experience swallowing issues following thyroidectomy. Although they are more frequently noticed in the late postoperative phase, these problems, characterized by delayed triggering and stasis of food, may be seen in the early postoperative period [21]. A study assessing a tool designed to calculate the likelihood of hypocalcemia following TT discovered that the decline in PTH levels 1 h and 8 h after surgery, in addition to magnesium levels on the second postoperative day, were reliable indicators of more severe hypocalcemia [22].

These results are in line with research showing that hypocalcemia is the most frequent side effect following thyroidectomy, with a frequency as high as 54.4%. Postoperative endocrinological symptoms, such as tingling, muscle cramps and fatigue, were reported in 12 (13.3%) patients. The most important factor influencing the occurrence of these symptoms was the degree of hypocalcemia ($p = 0.001$), while other factors such as the presence of goiter, hyperthyroidism and histopathological findings did not show a significant association. This confirms previous findings linking hypoparathyroidism with postoperative hypocalcemia [23]. Increased surgeon operative volume has been linked to better outcomes across various procedures, including thyroidectomy. A study showed that thyroidectomies on elderly persons were less common

among high-volume thyroid surgeons, who had fewer complications [24]. Risk-factor analysis showed that the presence of goiter alone significantly influences the occurrence of postoperative symptoms, which is consistent with the results of our study [25].

The decision regarding which surgical approach to take should be guided by the patient's risk factors, tumor characteristics and overall clinical context [26]. Thoracoscopy is considered the technique of choice for large retrosternal goiters descending into the mediastinum [27]. The treatment of autoimmune diseases of the thyroid gland such as Hashimoto's thyroiditis also requires surgical therapy. The solution would be improvement of diagnostics, good preoperative preparation, adequate intraoperative monitoring of the *n. laryngeus recurrens* and preoperative use of corticosteroids. Adequate and good results are directly related to precise surgical techniques, good operator experience and early voice rehabilitation [28].

CONCLUSION

Patients with VFP and hypocalcemia show the highest incidence of postoperative complications after TT due to HP-diagnosed goiter, follicular adenoma, and papillary thyroid carcinoma. These findings may be useful in identifying elderly patients at increased risk of complications after TT that can significantly affect their quality of life.

Conflict of interest: None declared.

REFERENCES

- Chahardahmasumi E, Salehidoost R, Amini M, Aminorroaya A, Rezvanian H, Kachooei A, et al. Assessment of the early and late complication after thyroidectomy. *Adv Biomed Res.* 2019;8:14. [DOI: 10.4103/abr.abr_3_19] [PMID: 30993084]
- El-Sebai Ali M, Atef Ebada H, El-Shaheed MA, Musaad AbdelFattah A, Kamal ES. Routine thyroidectomy with total laryngectomy: is it really indicated? A randomised controlled trial. *Ann Med Surg (Lond).* 2022;74:103309. [DOI: 10.1016/j.amsu.2022.103309] [PMID: 35145675]
- Sahli ZT, Zhou S, Sharma AK, Segev DL, Massie A, Zeiger MA, et al. Rising cost of thyroid surgery in adult patients. *J Surg Res.* 2021;260:28–37. [DOI: 10.1016/j.jss.2020.11.049] [PMID: 33316757]
- Bobanga ID, McHenry CR. Treatment of patients with Graves' disease and the appropriate extent of thyroidectomy. *Best Pract Res Clin Endocrinol Metab.* 2019;33(4):101319. [DOI: 10.1016/j.beem.2019.101319] [PMID: 31530446]
- Yahin E, Elbopa U, Kalender E. Regional reference values of thyroid gland volume in Turkish adults. *Srp Arh Celok Lek.* 2015;143(3–4):141–5. [DOI: 10.2298/SARH1504141S] [PMID: 26012121]
- Scerrino G, Melfa G, Raspanti C, Rotolo G, Salamone G, Licari L, et al. Minimally invasive video-assisted thyroidectomy: analysis of complications from a systematic review. *Surg Innov.* 2019;26(3):381–7. [DOI: 10.1177/1553350618823425] [PMID: 30632464]
- Metera A, Biancucci A, Natili A, Intini G, Graves CE. PTH after thyroidectomy as a predictor of post-operative hypocalcemia. *Diagnostics (Basel).* 2021;11(9):1733. [DOI: 10.3390/diagnostics11091733] [PMID: 34574074]
- Prduraru DN, Ion D, Carsote M, Andronic O, Bolocan A. Post-thyroidectomy hypocalcemia – risk factors and management. *Chirurgia (Bucur).* 2019;114(5):564–70. [DOI: 10.21614/chirurgia.114.5.564] [PMID: 31670631]
- Jin S, Sugitani I. Narrative review of management of thyroid surgery complications. *Gland Surg.* 2021;10(3):1135–46. [DOI: 10.21037/gs-20-859] [PMID: 33842257]
- Francis DO, Pearce EC, Ni S, Garrett CG, Penson DF. Epidemiology of vocal-fold paralyses after total thyroidectomy for well-differentiated thyroid cancer in a Medicare population. *Otolaryngol Head Neck Surg.* 2014;150(4):548–57. [DOI: 10.1177/0194599814521381] [PMID: 24482349]
- Pace-Asciak P, Russell JO, Tufano RP. Improving voice outcomes after thyroid surgery and ultrasound-guided ablation procedures. *Front Surg.* 2022;9:882594. [DOI: 10.3389/fsurg.2022.882594] [PMID: 35599805]
- Jan BS, Alamri AH, Alkaff HH, Almuqati WQ, Sayed SI, Abdelmonim SK, et al. Risk factors for postoperative hypocalcemia following total thyroidectomy: a retrospective study. *Ann Saudi Med.* 2024;44(1):39–47. [DOI: 10.5144/0256-4947.2024.39] [PMID: 38311865]
- Ryu CH, Lee SJ, Cho JG, Choi JJ, Choi YS, Hong YT, et al. Care and management of voice change in thyroid surgery: Korean Society of Laryngology, Phoniatrics and Logopedics clinical practice guideline. *Clin Exp Otorhinolaryngol.* 2022;15(1):24–48. [DOI: 10.21053/ceo.2021.00633] [PMID: 34098629]
- Gerardi I, Verro B, Amodei R, Richiusa P, Saraniti C. Thyroidectomy and its complications: a comprehensive analysis. *Biomedicines.* 2025;13(2):433. [DOI: 10.3390/biomedicines13020433] [PMID: 40002845]
- Chen HC, Pei YC, Fang TJ. Risk factors for thyroid surgery-related unilateral vocal-fold paralysis. *Laryngoscope.* 2019;129(1):275–83. [DOI: 10.1002/lary.27336] [PMID: 30284255]
- Kalezic N, Karadzic-Kocica M, Dimic N, Kocica M, Toškovac A, Jovanovic M, et al. Alcohol abuse as a risk factor for developing thyroid cancer. *Srp Arh Celok Lek.* 2021;149(3–4):185–8. [DOI: 10.2298/SARH201123113K]

17. Brown RL, de Souza JA, Cohen EE. Thyroid cancer: burden of illness and management of disease. *J Cancer*. 2011;2:193–9. [DOI: 10.7150/jca.2.193] [PMID: 21509149]
18. Stojanovic J, Milovanovic D, Mitrovic S, Jevtovic A, Bozovic N, Jovanovic M, et al. Videolaryngostroboscopy in early vocal-fold carcinoma diagnosis. *Srp Arh Celok Lek*. 2021;149(11–12):691–5. [DOI: 10.2298/SARH200529077S]
19. Huang CF, Jeng Y, Chen KD, Yu JK, Shih CM, Huang SM, et al. The preoperative evaluation prevents postoperative complications of thyroidectomy. *Ann Med Surg (Lond)*. 2014;4(1):5–10. [DOI: 10.1016/j.amsu.2014.11.005] [PMID: 25685337]
20. Gualniera P, Scurria S, Mondello C, Asmundo A, Sapienza D, Gianlorenzo D. Narrative review of proving the causal link of recurrent laryngeal-nerve injury and thyroidectomy: a medico-legal appraisal. *Gland Surg*. 2020;9(5):1564–72. [DOI: 10.21037/gs-20-203] [PMID: 33224832]
21. Elbeltagy YM, Bassiouny SE, Sobhy TS, Ismail AE, Teaima AA. Swallowing problems after thyroidectomy. *Int Arch Otorhinolaryngol*. 2021;26(3):e327–33. [DOI: 10.1055/s-0041-1730302] [PMID: 35846825]
22. Soares CSP, Koga KH, Moriguchi SM, Terra SA, Tagliarini JV, de Vasconcelos Affonso PHD, et al. Development of a tool to calculate the probability of hypocalcemia after total thyroidectomy: a prospective study. *Langenbecks Arch Surg*. 2024;409(1):33. [DOI: 10.1007/s00423-024-03229-8] [PMID: 38195723]
23. Suwannasarn M, Jongjaroenprasert W, Chayangsu P, Suvikapakornkul R, Sriphrapradang C. Single measurement of intact parathyroid hormone after thyroidectomy can predict transient and permanent hypoparathyroidism: a prospective study. *Asian J Surg*. 2017;40(5):350–6. [DOI: 10.1016/j.asjsur.2015.11.005] [PMID: 26780986]
24. Kim J, Seib CD. Operative management of thyroid disease in older adults. *J Endocr Soc*. 2023;7(7):bvad070. [DOI: 10.1210/jendso/bvad070] [PMID: 37324534]
25. Institute of Medicine (US) Committee on Standards for Developing Trustworthy Clinical Practice Guidelines; Graham R, Mancher M, Miller Wolman D, et al., editors. *Clinical practice guidelines we can trust*. Washington (DC): National Academies Press; 2011. [DOI: 10.17226/13058]
26. Javidi S, Sadrizadeh S, Sadrizadeh A, Bonakdaran S, Jarahi L. Postoperative complications and long-term outcomes after total and subtotal thyroidectomy: a retrospective study. *Sci Rep*. 2025;15:3705. [DOI: 10.1038/s41598-024-79860-8] [PMID: 39881145]
27. Jokic R, Antic J, Vorgucin I, Stajevic M, Nikin Z, Zeravica R, et al. Video-assisted thoracoscopic surgery for primary hyperparathyroidism with ectopic parathyroid adenoma in thymus. *Srp Arh Celok Lek*. 2021;149(5–6):365–9. [DOI: 10.2298/SARH200529042J]
28. Diklic A, Zivaljevic V, Paunovic I, Kalezic N, Tatic S. Surgical procedures in patients with thyroid autoimmune disease. *Srp Arh Celok Lek*. 2005;133(1):77–83. [DOI: 10.2298/SARH05S1077D] [PMID: 16405262]

Компликације после тоталне тиреоидектомије у корелацији са хистопатолошким и хормоналним налазима

Мирјана Ђатић^{1,2}, Далибор Врањеш^{1,2}, Вања Нежић², Бранко Јуришић^{1,2}, Ирма Овчина³

¹Универзитет у Бањој Луци, Медицински факултет, Бања Лука, Република Српска, Босна и Херцеговина;

²Универзитетски клинички центар Републике Српске, Клиника за болести уха, грла и носа, Бања Лука, Република Српска, Босна и Херцеговина;

³Универзитетски клинички центар Републике Српске, Клиника за унутрашње болести, Бања Лука, Република Српска, Босна и Херцеговина

САЖЕТАК

Увод/Циљ Тотална тиреоидектомија (ТТ) често се изводи у хирургији главе и врата и сматра се релативно једноставном процедуром, али је још увек праћена бројним компликацијама. Циљ ове студије био је да се испита улога и значај ТТ у настанку постоперативних компликација у корелацији са хистопатолошким (ХП) и ендокринолошким налазима.

Метод Студија пресека обухватила је 90 болесника који су били подвргнути операцији штитасте жлезде у Универзитетском клиничком центру Републике Српске у Бањој Луци, у Босни и Херцеговини, у периоду од 2022. до 2024. године. Преоперативно су урађене хормонске анализе, ултразвучни преглед штитасте жлезде и ендовидеоларингоскопија. Код свих болесника урађена је ТТ. Постоперативно су урађене ХП дијагностика, контролне хормонске анализе и ендовидеоларингоскопија.

Резултати Парализу гласница имало је 11 (12,2%) болесника. ХП анализа показала је да је у већини случајева узрок

био струма – код 37 (41,1%) болесника, затим фоликуларни аденом – код 25 (27,8%), папиларни карцином код 18 (20%) и остали узроци код 10 (11%) болесника. Од ендокринолошких компликација, хипокалцемија је била присутна код 67 (74,4%) болесника, а повишен ниво паратиroidног хормона код четири (4,4%) болесника.

Закључак Највећи број постоперативних компликација након ТТ јавља се код болесника са парализом гласница и хипокалцемијом због ХП дијагностиковане струме, фоликуларног аденома и папиларног карцинома штитасте жлезде. Ови налази могу бити корисни у идентификацији болесника старије животне доби с повећаним ризиком од компликација након ТТ, што може значајно утицати на квалитет њиховог живота.

Кључне речи: хипокалцемија; болести штитасте жлезде; парализа гласница