



ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

The amelioration of testicular ischemia-reperfusion injury in rats by oxymatrine

Si-Ming Wei^{1,2}, Yu-Min Huang³¹Zhejiang Shuren University, Shulan International Medical College, Hangzhou, Zhejiang, China;²Zhejiang Chinese Medical University, School of Nursing, Hangzhou, Zhejiang, China;³Zhejiang University, Hangzhou, College of Education, Department of Sports Science, Zhejiang, China**SUMMARY**

Introduction/Objective A crucial factor in testicular damage is the surplus of reactive oxygen species that arises from ischemia-reperfusion. Elevated levels of reactive oxygen species cause critical harm to cells. The present research focused on investigating the role of oxymatrine with antioxidative property in testicular ischemia-reperfusion injury induced by torsion and detorsion in a rat model.

Methods The study utilized a total of 60 rats, which were randomly assigned to one of three groups: a sham group with 20 rats, a testicular ischemia-reperfusion group with 20 rats, and a testicular ischemia-reperfusion group treated with oxymatrine, also consisting of 20 animals. Induction of testicular ischemia-reperfusion involved applying two hours of ischemia via 720° counterclockwise torsion to the left testis, with subsequent reperfusion initiated by detorsion. The activity of neutrophil nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, a catalyst for reactive oxygen species production, the concentration of a prevalent reactive oxygen species indicator malondialdehyde, and spermatogenesis were all detected in left testicular tissues collected at the four hours or three months post-detorsion.

Results Testicular ischemia-reperfusion notably elevated NADPH oxidase activity and malondialdehyde content, while it adversely affected spermatogenesis in the testes ($p < 0.001$). Conversely, the administration of oxymatrine significantly lowered both NADPH oxidase activity and malondialdehyde concentration, and it also improved spermatogenic function in the testes ($p < 0.001$).

Conclusion We found that oxymatrine attenuates injury from testicular ischemia-reperfusion, and this effect operates by limiting reactive oxygen species production through NADPH oxidase activity inhibition.

Keywords: oxymatrine; testicular torsion; ischemia-reperfusion

INTRODUCTION

The occurrence of testicular torsion spans all ages, yet it predominantly affects male infants and teenagers in the 12–18 age range, making it a standard emergency seen in urological practice [1]. Testicular torsion occurs when the spermatic cord becomes twisted, leading to a partial or total interruption of blood supply to the testicle. Early detection and quick surgical untwisting must be performed to reinstate testicular blood flow and forestall ischemic necrosis. Although the twisted testicle is surgically untwisted, subsequent atrophy and loss of function afflict 13.5–68% of such cases [2]. Ischemia-reperfusion injury remains the most recognized cause of testicular damage subsequent to testicular torsion and its surgical relief [3]. A crucial factor in testicular damage is the surplus of reactive oxygen species that arises from ischemia-reperfusion [3]. Elevated levels of reactive oxygen species (e.g., hydroxyl radical, superoxide anion, hydrogen peroxide) cause critical harm to cells by oxidizing membrane lipids, altering protein structures, and breaking DNA strands [3].

As of now, patients have no access to clinically utilized drugs that can reduce testicular ischemia-reperfusion injury. Oxymatrine is a type of quinolizidine alkaloid mainly isolated

from *Sophora flavescens*, a plant utilized in traditional Chinese herbal practice [4, 5, 6]. $C_{15}H_{24}N_2O_2$ represents its molecular formula, while its weight is calculated as 264.36 g/mol [7, 8]. The antioxidative and anti-inflammatory properties of oxymatrine are among the numerous bioactivities that have been identified through multiple investigations [9, 10, 11]. Studies further report that oxymatrine mitigates damage from ischemia-reperfusion in several areas: the lung, brain, intestine, kidney, and liver [12–17]. Despite this, research has not addressed the therapeutic effect of oxymatrine on testicular ischemia-reperfusion injury. The present research focused on investigating the role of oxymatrine in testicular ischemia-reperfusion injury induced by torsion and detorsion in a rat model.

METHODS**Study subjects**

The animals in this study consisted of sixty male Sprague-Dawley rats weighing between 250 and 300 grams, sourced from Shanghai Laboratory Animal Center company, located in Shanghai, China. Their housing consisted of an air-conditioned room with tightly regulated

Received • Примљено:
February 4, 2026

Revised • Ревизија:
July 13, 2026

Accepted • Прихваћено:
July 15, 2026

Online first: July 30, 2026

Correspondence to:

Si-Ming WEI
No. 8 Shuren Street
310015 Hangzhou
Zhejiang
China
wsm1971@hotmail.com

conditions: a temperature of 21°C, varying by no more than 1°C, a relative humidity of 55%, varying within 5%, and a standard 12-hour light followed by 12-hour dark schedule. Throughout the experiment, the rats could consume standard food and water at will. The Zhejiang Shuren University Ethics Committee provided ethical approval for all study protocols, which bears the reference number 202309021. The research methods involving animals were designed and executed following universally accepted ethical rules.

Medicinal compound and experimental substances

Sigma-Aldrich (Sigma-Aldrich, St. Louis, MO, USA) was the provider of ketamine, oxymatrine, hematoxylin, and eosin. The Jiancheng Bioengineering Institute (Nanjing, China) supplied the kits for analyzing nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and malondialdehyde. The remaining reagents utilized during the investigation were premium grade and sourced from the market.

Research plan

A random distribution was performed for 60 Sprague-Dawley rats among three groups, namely sham, testicular ischemia-reperfusion, and oxymatrine-treated groups, with 20 animals in each group. The rats received ketamine (50 mg/kg) intraperitoneally for anesthesia and were then positioned on the operating table. Sterility was maintained throughout all experimental manipulations. The left testis was retrieved by a left ilioinguinal surgical approach. To simulate the surgical procedure, an 11-0 atraumatic silk suture was positioned within the tunica albuginea of animals belonging to the sham group [18]. The procedure concluded with the replacement of the testis into the scrotum and the suturing of the skin. Ischemia in the testicular ischemia-reperfusion model was produced through a 720° counterclockwise twist applied to the left testis [18]. An 11-0 silk suture was used to attach the torsioned testis to the scrotum, thereby preserving the ischemic state. At the conclusion of the two-hour interval, restoring the testis to its natural alignment through untwisting enabled reperfusion. In the experimental group receiving oxymatrine, a two-hour period of testicular ischemia preceded reperfusion, at which time a 40 mg/kg dose of oxymatrine was injected intraperitoneally. The justification for the oxymatrine dose employed originated from various prior studies [15, 16, 17]. The rats received nalbuphine (2 mg/kg) subcutaneously for postoperative analgesia. Reperfusion was conducted for four hours, after which 10 rats in each group underwent left orchietomy with the goal of assessing NADPH oxidase activity and malondialdehyde concentration. The remaining ten animals in each group were subjected to left orchietomy after three months of reperfusion, allowing for the measurement of testicular spermatogenesis. Once the experimental procedures ended, all animals received a terminal intraperitoneal dose of pentobarbital sodium (200 mg/kg) to ensure a painless death.

Testicular tissue analysis for NADPH oxidase activity

After being homogenized in lysis buffer kept on ice, the samples were centrifuged at 4°C and 10,000 × g for a duration of 10 minutes. To measure NADPH oxidase activity, the supernatant was harvested with care. Assessment of NADPH oxidase activity relied on a kit purchased from a commercial source.

Determining malondialdehyde concentration in tissue from the testis

On ice, testicular tissue underwent homogenization in a buffer designed for lysis with malondialdehyde. The supernatant intended for the malondialdehyde assay was acquired by centrifuging the tissue homogenate at 5000 g for a duration of 15 minutes under 4°C conditions. Measurement of malondialdehyde levels in testicular tissue was performed via the thiobarbituric acid reactive substance assay, following the protocol supplied with a commercial kit.

Evaluation of sperm production within the testicles

Parameters such as seminiferous tubule epithelial layer number, Johnsen score, testicular weight, and seminiferous tubular diameter served as indicators for assessing testicular spermatogenesis [19]. After excision, the testis was placed on a scale for weight measurement. Histopathological examination was facilitated by fixing part of the testis in Bouin's fluid, dehydrating it with increasing ethanol concentrations, and embedding it in paraffin. The paraffin block was sectioned on a rotary microtome to produce slices of 5 µm in thickness. Staining with hematoxylin and eosin was applied to the sections once the routine processes of deparaffinization and hydration were complete. Light microscopic evaluation of the testicular sections was performed by an assessor who was not informed of their origin. The evaluation procedure consisted of selecting 20 seminiferous tubules with circular or near-circular profiles from each histological section and measuring their epithelial layer number, Johnsen score, and diameter. To assess the seminiferous tubule epithelial layers, counts were performed starting at the basement membrane and proceeding inward to the lumen. Histology of the seminiferous tubules was scored in accordance with Johnsen's system, which grades findings based on germinal epithelial maturation [20]. Seminiferous tubules were graded according to Johnsen's criteria, with scores spanning from 1 to 10 [20]. A score of 1 is assigned when the seminiferous tubule is entirely devoid of cells. The highest score of 10 is awarded for complete spermatogenesis accompanied by plentiful spermatozoa, a germinal epithelium of standard thickness, and a patent tubular lumen. Using a microscope that had an ocular micrometer, researcher estimated the diameter of the seminiferous tubules.

Statistical analysis

The GraphPad Prism 4.0 (GraphPad Software, Boston, MA, USA) served as the tool for statistical analysis. Using the Shapiro–Wilk test, we confirmed that the data exhibited a normal distribution. The variance homogeneity was verified by Bartlett's test. Data from the experiments were summarized using means and standard deviations. For intergroup comparisons, we employed one-way analysis of variance and subsequently conducted the Student–Newman–Keuls multiple comparison test. A *p*-value lower than 0.05 indicated that the differences were statistically significant.

Ethics: The study was approved by the Ethics Committee of the Zhejiang Shuren University (No.:202309021).

RESULTS

NADPH oxidase activity observed in testes

The activity of NADPH oxidase in testes was significantly higher in the ischemia-reperfusion group than in the sham group, a result depicted in Figure 1 ($p < 0.001$). Following testicular ischemia-reperfusion, the observed upregulation of NADPH oxidase activity was significantly decreased by oxymatrine treatment ($p < 0.001$).

Testicular tissue levels of malondialdehyde

Figure 2 illustrated a significant elevation in testicular malondialdehyde levels in the ischemia-reperfusion group ($p < 0.001$) relative to the sham-operated controls. Relative to animals subjected to ischemia-reperfusion, those receiving oxymatrine showed significantly ($p < 0.001$) reduced malondialdehyde levels in testicular tissue.

Evidence of spermatogenic activity in testicular tissue

Testicular ischemia-reperfusion injury led to marked declines ($p < 0.001$) in several histological and morphometric parameters, including number of epithelial layers within the tubules, Johnsen scoring, weight of the testes, and diameter of the seminiferous tubules, relative to sham controls (Figures 3 and 4). Oxymatrine-treated animals exhibited significantly ($p < 0.001$) better outcomes for all four testicular parameters compared to those subjected to ischemia-reperfusion injury.

DISCUSSION

In the population of males under 25 years old, testicular torsion arises at a rate of one per 4000 [2]. For the conservation of testicular fertility, early detection and prompt surgical correction of torsion are essential. Testicular infarction occurs if torsion of the testis continues beyond six hours [3]. Timely surgery within a five-hour window




Figure 1. Testicular nicotinamide adenine dinucleotide phosphate (NADPH) oxidase activity was analyzed in the following conditions: sham operation, testicular ischemia-reperfusion (tIR), and oxymatrine (OM) treatment; data ($n = 10$) from the experiments were summarized using means and standard deviations; a – with a *p*-value of less than 0.001 vs. the sham control; b: – with a *p*-value of less than 0.001 vs. the tIR group




Figure 2. Testicular malondialdehyde level was analyzed in the following conditions: sham operation, testicular ischemia-reperfusion (tIR), and oxymatrine (OM) treatment; data ($n = 10$) from the experiments were summarized using means and standard deviations; a: with a *p*-value of less than 0.001 vs. the sham control; b: with a *p*-value of less than 0.01 vs. the sham control; c: with a *p*-value of less than 0.001 vs. the tIR group

of symptom appearance fails to avert testicular atrophy in 27% of affected individuals [21]. The testes in our investigation remained viable after undergoing torsion for two hours and then being surgically restored. Severe impairment of spermatogenesis was observed in the testis three months post-detorsion, a consequence of the initial torsion-detorsion event. Key indicators of this damage, detailed in Figures 3 and 4, included a significantly reduced epithelial layer count in seminiferous tubules, a lower Johnsen score, diminished testicular weight, and narrower seminiferous tubular diameters.

From a pathophysiological perspective, torsion and detorsion of the testis is a quintessential ischemia-reperfusion phenomenon. Reactive oxygen species are produced in excess during ischemia and reperfusion of the testis [22]. The oxidation of proteins, DNA, and membrane lipids by surplus reactive oxygen species is a mechanism capable of initiating cellular damage [3]. The extreme reactivity and fleeting existence of reactive oxygen species make them exceedingly difficult to quantify in a direct manner [18]. Malondialdehyde, produced steadily at the end of lipid




Figure 3. (A) – number of epithelial layers within the tubules; (B) – Johnsen scoring; (C) – weight of the testes; (D) – diameter of the seminiferous tubules were analyzed in the following conditions: sham operation, testicular ischemia-reperfusion (tIR), and oxymatrine (OM) treatment; data (n = 10) from the experiments were summarized using means and standard deviations; a – with a p-value of less than 0.001 vs. the sham control; b – with a p-value of less than 0.001 vs. the tIR group; c – with a p-value of less than 0.05 vs. the sham control; d – with a p-value of less than 0.01 vs. the sham control

hepatitis B, plaque psoriasis, and liver fibrosis from chronic viral hepatitis [23–27]. Therefore, the findings provide a foundation for the potential therapeutic application of oxymatrine in cases of testicular ischemia-reperfusion insult. Yet, the pathways responsible for the decrease in reactive oxygen species induced by oxymatrine are still poorly characterized.

Infiltrating neutrophils represent a primary cause of excessive reactive oxygen species production in tissues undergoing ischemia-reperfusion [28]. In neutrophils, NADPH oxidase is a key contributor to the formation of reactive oxygen species [28]. Ischemia promotes calcium influx into neutrophils, which in turn activates NADPH oxidase [29]. The reperfusion phase is characterized by the restoration of oxygen to previously ischemic tissue. Superoxide anion is generated from oxygen through catalysis by activated NADPH oxidase




Figure 4. Testis tissue sections were illustrated for the three experimental conditions: sham operation, testicular ischemia-reperfusion, and oxymatrine treatment; at a magnification of $\times 200$, testicular tissue sections underwent microscopic evaluation after being stained with hematoxylin-eosin; (A) – microscopic assessment of sham-operated testes indicated regular layering of the epithelium, full spermatogenic development ranging from early germ cells to spermatozoa (indicated by arrows), and standard tubular diameter; every seminiferous tubule contained a clearly open central lumen; (B) – the ischemia-reperfusion group presented with testicular structural anomalies: seminiferous tubules were atrophic (arrow), their epithelial layers were diminished (arrowhead), and germ cells showed maturation arrest (arrowhead); (C) – the gross morphology of the testes in the oxymatrine-treated group closely matched that of the sham group; inside the tubules, researchers noted numerous mature sperm cells, which are pointed out with an arrow; an arrowhead indicates sloughed germinal cells within the tubular lumen, a condition that often led to blockage; 40 micrometers are the length represented by the scale bar

peroxidation initiated by reactive oxygen species, is largely considered a sensitive sign of their presence and activity [19]. We report here that testicular malondialdehyde levels rose significantly while spermatogenesis was significantly suppressed following ischemia-reperfusion (Figures 2, 3, and 4). The evidence points to overproduction of reactive oxygen species as the underlying cause of testicular ischemia-reperfusion injury. The study demonstrated that oxymatrine supplementation lowered malondialdehyde concentration while boosting testicular spermatogenesis (Figures 2, 3, and 4). According to our results, the antioxidant capacity of oxymatrine underlies its ability to mitigate testicular ischemia-reperfusion injury. Furthermore, the safety and therapeutic effectiveness of oxymatrine have been confirmed in clinical settings for treating chronic

[29]. Superoxide anions undergo a self-reaction to generate hydrogen peroxide [30]. The interaction of superoxide anion and hydrogen peroxide yields hydroxyl radical [19]. Thus, reactive oxygen species accumulate in greater amounts following ischemia and subsequent reperfusion. Testicular ischemia-reperfusion in this study produced substantial rises in NADPH oxidase activity and malondialdehyde content, coupled with a pronounced decrease in spermatogenesis within the testes, as illustrated in Figures 1, 2, 3, and 4. The study implies that spermatogenic injury arises from reactive oxygen species overproduction, which itself results from upregulated NADPH oxidase activity in neutrophils following testicular ischemia-reperfusion. Moreover, our results revealed that oxymatrine significantly suppressed NADPH oxidase activity and reduced

malondialdehyde content, while effectively restoring spermatogenesis in testicular tissue (Figures 1, 2, 3, and 4). The data imply that decreased reactive oxygen species production achieved by inhibiting NADPH oxidase activity in neutrophils underlies the protective role of oxymatrine in testicular spermatogenesis.

Khorsand et al. [3] evaluated whether naringin, which is a flavanone glycoside possessing well-documented antioxidant, anti-inflammatory, and anti-apoptotic effects, could prevent testicular ischemia/reperfusion damage induced by torsion/detorsion in mice [3]. Compared to controls, the torsion/detorsion animals exhibited a significant reduction in sperm quality based on concentration, motility, and kinematics, a depletion of antioxidant reserves such as superoxide dismutase, total antioxidant capacity, and glutathione peroxidase, a drop in hormones like follicle-stimulating hormone, testosterone, and luteinizing hormone, a rise in malondialdehyde and apoptosis signals Bax, Bcl-2, and caspase-3, and a loss of fertility. Naringin produced a dose-dependent improvement in sperm indices, antioxidant defense, testicular structure, endocrine parameters, and fertility, particularly when given at 100 or 200 mg/kg. At 50 mg/kg, the treatment exhibited weak effectiveness. Naringin exerted its apoptosis-inhibiting effect via the enhancement of Bcl-2 and the simultaneous reduction of Bax and caspase-3. These findings suggest that testicular ischemia/reperfusion injury is significantly alleviated by naringin, with the 100 to 200 mg/kg doses being especially effective. The comparison reveals that, aside from malondialdehyde and testicular histology, the experimental design of Khorsand et al. [3] was distinct from ours with respect to the other assessed variables and the therapeutic strategies. The next phase of this research will integrate measurements of the other assessed factors,

and evaluate the relative potency of oxymatrine vs. naringin for treating testicular ischemia-reperfusion damage, identifying the optimal therapeutic candidate.

Several limitations within our research must be thoroughly explored by future studies. Firstly, the impact of unilateral testicular torsion on the contralateral testicle was left unexamined. Secondly, our research only used one oxymatrine dose and lacked dose-response evaluation. Thirdly, our study only used a single animal species and lacked molecular confirmation of the proposed mechanism (e.g., NADPH oxidase isoforms, inflammatory cytokines, antioxidant enzymes, Bax, Bcl-2, and caspase-3).

CONCLUSION

This study presents the first indication that the protection offered by oxymatrine against ischemia/reperfusion-induced testicular injury involves the inhibition of neutrophil NADPH oxidase activity, resulting in reduced generation of reactive oxygen species. According to the results, oxymatrine may function as a viable therapeutic option in cases of testicular torsion-detorsion. The potential clinical usefulness of oxymatrine awaits confirmation from trials conducted in humans.

ACKNOWLEDGMENT

Financial support for this research was provided through Grant LY19H040001 from the Zhejiang Provincial Natural Science Foundation in China.

Conflict of interest: None declared.

REFERENCES

- Shields LB, Jawad K, Rosenberg E. Higher orchiectomy rates in pre-pubescent testicular torsion: A call for earlier recognition and heightened awareness. *Cureus*. 2025;17(9): e93512. [DOI: 10.7759/cureus.93512] [PMID: 41170223]
- Refaie MMM, Abdel-Hakeem EA, Ahmed Ibrahim R, Hassan Mohamed Z, Shehata S, Soltan Shrekawy El Bakry H, et al. Role of sacubitril/valsartan in testicular ischemia reperfusion injury via modulation of AngII/ASK/NF- κ B and JAK/STAT pathways in juvenile rats. *Br J Pharmacol*. 2025;182(18):4343–62. [DOI: 10.1111/bph.70092] [PMID: 40456625]
- Khorsand K, Najafpour A, Valilou M, Soleimanzadeh A. Naringin attenuates testicular ischemia/reperfusion injury in a mouse torsion/detorsion model via intraperitoneal administration. *Pediatr Surg Int*. 2026;42(1):59. [DOI: 10.1007/s00383-025-06286-2] [PMID: 41492007]
- Liu Y, Wang H, Liu N, Du J, Lan X, Qi X, et al. Oxymatrine protects neonatal rat against hypoxic-ischemic brain damage via PI3K/Akt/GSK3 β pathway. *Life Sci*. 2020; 254:116444. [DOI: 10.1016/j.lfs.2019.04.070] [PMID: 31102745]
- Xiong Z, Xu J, Liu X. Oxymatrine exerts a protective effect in myocardial ischemia/reperfusion-induced acute lung injury by inhibiting autophagy in diabetic rats. *Mol Med Rep*. 2021;23(3):183. [DOI: 10.3892/mmr.2021.11822] [PMID: 33398371]
- Huang Y, Long X, Li X, Li S, He J. The Role of oxymatrine in amelioration of acute lung injury subjected to myocardial I/R by inhibiting endoplasmic reticulum stress in diabetic rats. *Evid Based Complement Alternat Med*. 2020;2020:8836904. [DOI: 10.1155/2020/8836904] [PMID: 33293996]
- Wang WJ, Ma YM, He MT, Zhang DH, Wang R, Jing L, et al. Oxymatrine alleviates hyperglycemic cerebral ischemia/reperfusion injury via protecting microvessel. *Neurochem Res*. 2022;47(5):1369–82. [DOI: 10.1007/s11064-022-03535-x] [PMID: 35099719]
- Jiang G, Liu X, Wang M, Chen H, Chen Z, Qiu T. Oxymatrine ameliorates renal ischemia-reperfusion injury from oxidative stress through Nrf2/HO-1 pathway. *Acta Cir Bras*. 2015;30(6):422–9. [DOI: 10.1590/S0102-865020150060000008] [PMID: 26108031]
- Zhou S, Qiao B, Chu X, Kong Q. Oxymatrine attenuates cognitive deficits through SIRT1-mediated autophagy in ischemic stroke. *J Neuroimmunol*. 2018;323:136–42. [DOI: 10.1016/j.jneuroim.2018.06.018] [PMID: 30196826]
- Zhu B, Yang JR, Chen SF, Jiang YQ. The attenuation of lung ischemia reperfusion injury by oxymatrine. *Cell Biochem Biophys*. 2014;70(1):333–6. [DOI: 10.1007/s12013-014-9917-4] [PMID: 24696074]
- Jiao-Yan Y, Qing-Qing L, Xi L, Mei Z, Ting S, Na H, et al. Oxymatrine improves blood-brain barrier integrity after cerebral ischemia-reperfusion injury by downregulating CAV1 and MMP9 expression. *Phytomedicine*. 2021; 84:153505. [DOI: 10.1016/j.phymed.2021.153505] [PMID: 33626426]
- Zhu B, Yang J, Chen S, Zhang P, Shen L, Li X, et al. Oxymatrine on Hsp90a expression and apoptosis in a model of lung ischemia-reperfusion injury. *Exp Ther Med*. 2017;13(4):1381–5. [DOI: 10.3892/etm.2017.4098] [PMID: 28413481]
- Ma CS, Han B, Liu YX, Shi CK, Li DL, Guo JF, et al. Oxymatrine alleviates cerebral ischemia/reperfusion injury by targeting HDAC1 to regulate mitochondria-related autophagy and oxidative stress. *Mol Neurobiol*. 2025;63(1):219. [DOI: 10.1007/s12035-025-05423-1] [PMID: 41317220]

14. Li M, Zhang X, Cui L, Yang R, Wang L, Liu L, et al. The neuroprotection of oxymatrine in cerebral ischemia/reperfusion is related to nuclear factor erythroid 2-related factor 2 (nrf2)-mediated antioxidant response: role of nrf2 and hemeoxygenase-1 expression. *Biol Pharm Bull.* 2011;34(5):595–601. [DOI: 10.1248/bpb.34.595] [PMID: 21532144]
15. Zhao J, Yu S, Tong L, Zhang F, Jiang X, Pan S, et al. Oxymatrine attenuates intestinal ischemia/reperfusion injury in rats. *Surg Today.* 2008;38(10):931–7. [DOI: 10.1007/s00595-008-3785-8] [PMID: 18820869]
16. Ozturk H, Cetinkaya A, Yilmaz F, Ozturk H. Protective effect of oxymatrine against renal ischemia/reperfusion injury in rats. *Bratisl Lek Listy.* 2017;118(4):217–22. [DOI: 10.4149/BLL_2017_043] [PMID: 28471232]
17. Jiang H, Meng F, Li J, Sun X. Anti-apoptosis effects of oxymatrine protect the liver from warm ischemia reperfusion injury in rats. *World J Surg.* 2005;29(11):1397–401. [DOI: 10.1007/s00268-005-7885-y] [PMID: 16222447]
18. Wei SM, Huang YM. Safranin ameliorates testicular ischemia-reperfusion injury in testicular torsion-detorsion rat model. *Rev Int Androl.* 2024;22(4):33–41. [DOI: 10.22514/j.androl.2024.028] [PMID: 39748768]
19. Wei SM, Huang YM. Protective effect of allicin on ischemia-reperfusion injury caused by testicular torsion-detorsion in rats. *Eur Rev Med Pharmacol Sci.* 2024;28(7):2817–26. [DOI: 10.26355/eurrev_202404_35910] [PMID: 38639521]
20. Johnsen SG. Testicular biopsy score count—a method for registration of spermatogenesis in human testes: normal values and results in 335 hypogonadal males. *Hormones.* 1970;1(1):2–25. [DOI: 10.1159/000178170] [PMID: 5527187]
21. Sessions AE, Rabinowitz R, Hulbert WC, Goldstein MM, Mevorach RA. Testicular torsion: direction, degree, duration and disinformation. *J Urol.* 2003;169(2):663–5. [DOI: 10.1097/01.ju.0000047381.36380.0e] [PMID: 12544339]
22. Wei SM, Huang YM. Effect of sulforaphane on testicular ischemia-reperfusion injury induced by testicular torsion-detorsion in rats. *Sci Rep.* 2024;14(1):23420. [DOI: 10.1038/s41598-024-74756-z] [PMID: 39379457]
23. Lu LG, Zeng MD, Mao YM, Li JQ, Wan MB, Li CZ, et al. Oxymatrine therapy for chronic hepatitis B: a randomized double-blind and placebo-controlled multi-center trial. *World J Gastroenterol.* 2003;9(11):2480–3. [DOI: 10.3748/wjg.v9.i11.2480] [PMID: 14606080]
24. Gu XB, Yang XJ, Hua Z, Lu ZH, Zhang B, Zhu YF, et al. Effect of oxymatrine on specific cytotoxic T lymphocyte surface programmed death receptor-1 expression in patients with chronic hepatitis B. *Chin Med J (Engl).* 2012;125(8):1434–8. [DOI: 10.3760/cma.j.issn.0366-6999.2012.08.014] [PMID: 22613649]
25. Zhou H, Shi HJ, Yang J, Chen WG, Xia L, Song HB, et al. Efficacy of oxymatrine for treatment and relapse suppression of severe plaque psoriasis: results from a single-blinded randomized controlled clinical trial. *Br J Dermatol.* 2017;176(6):1446–55. [DOI: 10.1111/bjd.15316] [PMID: 28112799]
26. Shi HJ, Zhou H, Ma AL, Wang L, Gao Q, Zhang N, et al. Oxymatrine therapy inhibited epidermal cell proliferation and apoptosis in severe plaque psoriasis. *Br J Dermatol.* 2019;181(5):1028–37. [DOI: 10.1111/bjd.17852] [PMID: 30822359]
27. Mao YM, Zeng MD, Lu LG, Wan MB, Li CZ, Chen CW, et al. Capsule oxymatrine in treatment of hepatic fibrosis due to chronic viral hepatitis: a randomized, double blind, placebo-controlled, multicenter clinical study. *World J Gastroenterol.* 2004;10(22):3269–73. [DOI: 10.3748/wjg.v10.i22.3269] [PMID: 15484298]
28. Ogunleye OD, Afolabi OA, Saka WA, Olusola BO, Ajike RA, Oladokun OO, et al. Possible role of vitamins C and E co-administration in the prevention of testicular ischemia-reperfusion injury following surgical repair of torsion of the testis. *Front Nutr.* 2025; 12:1660240. [DOI: 10.3389/fnut.2025.1660240] [PMID: 41256933]
29. Kračun D, Görlach A, Snedeker JG, Buschmann J. Reactive oxygen species in tendon injury and repair. *Redox Biol.* 2025; 81:103568. [DOI: 10.1016/j.redox.2025.103568] [PMID: 40023978]
30. Lee J, Yeo SG, Lee JM, Kim SS, Lee JW, Chung N, et al. Expression of free radicals and reactive oxygen species in endometriosis: current knowledge and its implications. *Antioxidants (Basel).* 2025;14(7):877. [DOI: 10.3390/antiox14070877] [PMID: 40722981]

Ублажавање повреда тестиса услед исхемије-реперфузије код пацова оксиматриному

Си-Минг Беј^{1,2}, Ју-Мин Хуанг³

¹Универзитет Жеђанг Шужен, Међународни медицински колеџ Шулан, Ханџоу, провинција Жеђанг, Кина;

²Универзитет традиционалне кинеске медицине Жеђанг, Факултет за сестринство, Ханџоу, провинција Жеђанг, Кина;

³Универзитет Жеђанг, Факултет за образовање, Катедра за спортске науке, Ханџоу, провинција Жеђанг, Кина

САЖЕТАК

Увод/Циљ Кључни фактор оштећења тестиса је вишак реактивних врста кисеоника који настаје услед исхемије-реперфузије. Повећани нивои реактивних врста кисеоника наносе озбиљну штету ћелијама. Ово истраживање било је усмерено на испитивање улоге оксиматрина са антиоксидативним својствима у исхемијско-реперфузијској повреди тестиса изазваној торзијом и деторзијом у моделу пацова.

Метод У студији је коришћено укупно 60 пацова, који су насумично распоређени у једну од три групе: лажно оперисана група са 20 пацова, група са исхемијом-реперфузијом тестиса са 20 пацова и група са исхемијом-реперфузијом тестиса третирана оксиматрином, која се такође састојала од 20 животиња. Индукција исхемије-реперфузије тестиса подразумевала је двочасовну исхемију путем торзије левог тестиса за 720° у смеру супротном од казаљке на сату, а накнадна реперфузија покренута је деторзијом. Активност неутрофилне никотинамид-аденин-динуклеотид-фосфатне

(NADPH) оксидазе, катализатора за производњу реактивних врста кисеоника, концентрација малондиалдехида, показатеља реактивних врста кисеоника, и сперматогенеза испитивани су у ткивима левог тестиса прикупљеним четири сата или три месеца после деторзије.

Резултати Тестикуларна исхемија-реперфузија значајно је повећала активност NADPH оксидазе и садржај малондиалдехида, док је негативно утицала на сперматогенезу у тестисима ($p < 0,001$). Супротно томе, примена оксиматрина значајно је смањила активност NADPH оксидазе и концентрацију малондиалдехида, а такође је побољшала сперматогену функцију тестиса ($p < 0,001$).

Закључак Утврђено је да оксиматрин ублажава повреде настале услед исхемије-реперфузије тестиса, а овај ефекат се остварује ограничавањем производње реактивних врста кисеоника инхибицијом активности NADPH оксидазе.

Кључне речи: оксиматрин; торзија тестиса; исхемија-реперфузија