



Paper Accepted^{*}

ISSN Online 2406-0895

Original Article / Оригинални рад

Jelena B. Ilić-Živojinović^{1†}, Branislav B. Ilić², Dušan Backović¹, Milena Tomanić¹, Aleksandar Gavrilović³, Ljiljana Bogdanović⁴

Knowledge and attitudes on medical waste management among Belgrade medical and dental students

Знање и ставови београдских студената медицине и стоматологије о управљању медицинским отпадом

¹University of Belgrade, Faculty of Medicine, Institute of Hygiene and Medical Ecology, Belgrade, Serbia;
 ²University of Belgrade, School of Dental Medicine, Clinic for Oral Surgery, Belgrade, Serbia;
 ³University of Kragujevac, Faculty of Medical Sciences, Department of Neurology, Kragujevac, Serbia;
 ⁴University of Belgrade, Institute of Pathology, Faculty of Medicine, Belgrade, Serbia;

Received: April 5, 2018 Revised: July 26, 2018 Accepted: October 4, 2018 Online First: October 16, 2018 DOI: https://doi.org/10.2298/SARH180405065I

[†]**Correspondence to:** Jelena ILIĆ-ŽIVOJINOVIĆ, Institute of Hygiene and Medical Ecology, Faculty of Medicine, University of Belgrade, Dr Subotića 8 11000 Belgrade, Serbia Email: **jelenil77@gmail.com**

^{*} Accepted papers are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. Srp Arh Celok Lek. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

Knowledge and attitudes on medical waste management among Belgrade medical and dental students

Знање и ставови београдских студената медицине и стоматологије о управљању медицинским отпадом

SUMMARY

Introduction/Objective Knowledge and practical skills in medical waste (MW) management are of equal importance for medical and dental doctors. The first comparative study on knowledge and skills in the field of medical waste management among Belgrade students, was conducted with the goal examining whether this extremely important area is governed equally well by students of medicine and dentistry.

Methods A cross-sectional study included 558 students of the sixth year of studies (430 medical and 128 dental students) which completed an anonymous semi-structured questionnaire to determine attitudes and knowledge on MW management.

Results The majority of medical and dental students had no training in MW management (79.5% and 74.6%, respectively). Dental students use protective equipment more frequently than medical students (94.5% vs. 42.0%, p < 0.001). However, full vaccinal protection against hepatitis B is better among medical students compared to dental students (57.7% vs. 39.1%, p <0.001). Complete knowledge on postexposal prophylaxis is better among medical students compared to dental students (44.5% vs. 13.3%, p <0.001). However, dental students are more disciplined in reporting injuries (63.1% vs. 52.4%, p = 0.038). The students' knowledge on primary separation of infectious waste and used needles is better among dental students compared to medical students /correct answers, 93.0% vs. 77.8% (p<0.001) and 80.3% vs. 70.4% (p=0.007), respectively.

Conclusion Dental students show better knowledge on MW management and are more disciplined in using personal infection protection compared to medical students. The students support continuing training on MW management and investigations on this topic.

Keywords: medical waste; safety; education; medical students; dental students

Сажетак

Увод/Циљ Знање и практичне вештине управљања медицинским отпадом су од посебног значаја за докторе медицине и стоматологије. Компаративна студија о знању и вештинама из области управљања медицинским отпадом спроведена је међу београдским студентима први пут, у циљу испитивања да ли овом изузетно важном облашћу студенти медицине и стоматологије владају подједнако добро.

2

Методе Ова студија пресека обухватила је 558 студената шесте године студија на Београдском Универзитету (430 студената медицине и 128 студената стоматологије) који су попунили анонимни упитник о знању и ставовима о управљању медицинским отпадом.

Резултати Већина студената стоматологије и медицине нису имали никакву посебну обуку из управљања медицинским отпадом (79,5% и Студенти стоматологије су чешће 74,6%). користили заштитну опрему (94,5% према 42.0%, р < 0.001). Вакцинална заштита од хепатитиса Б је комплетнија међу студентима медицине у односу на студенте стоматологије (57,7% према 39,1% *p*<0.001). Знање о профилакси после изложености боља је међу студентима медицине (44,5% према <0.001). 13,3%, р Међутим, студенти стоматологије ажурнији погледу cy y пријављивања повреда на радном месту (63,1% према 52,4%, *p* = 0.038). Знање студената о примарној сепарацији инфективног отпада и коришћених игала је боље међу будућим стоматолозима (тачни одговори 93,0% насупрот 77,8%; *p*<0.001 и 80,3% према 70,4%; *p*=0.007).

Закључак Студенти стоматологије имају боље знање о управљању медицинским отпадом и дисциплинованији су у погледу коришћења заштитне опреме на раду у односу на студенте медицине. Студенти подржавају континуирану едукацију о управљању медицинским отпадом и даља истраживања о овој теми.

Кључне речи: медицински отпад; сигурност; едукација; студенти медицине; студенти стоматологије

INTRODUCTION

The term medical waste (MW) refers to all the waste generated within health-care facilities, research centers and laboratories. The term MW consists of materials ranging from used needles to body parts, diagnostic samples, blood, chemicals, pharmaceuticals, and radioactive materials. From 10% to 25% of all MW is hazardous and may cause a variety of environmental and health risks [1,2].

A quarter of all MW in Serbia is hazardous [3] and infectious MW is the largest part of it [4]. The annual production of infectious waste in Serbia is between 4500 and 5000 tons [5]. Since 2006 a national system for safe MW management has been developed and all infectious MW has been sterilized. Serbia has reduced the amount of hazardous MW by 50% by introducing a waste separation process in health-care facilities [6].

Recognizing the importance of knowledge on MW management for health-care professionals, Serbian medical faculties have recently introduced MW topic into the curriculum on the final year of their practice. To assess the quality of undergraduate education on MW management it is important to check the students' retention of knowledge. Further, although the knowledge in this field is of equal importance for medical and dental doctors, it is not clear whether medical and dental students adopt this necessary knowledge equally. For these reasons, we undertake this comparative study on knowledge and attitudes on MW management among Belgrade medical and dental students.

METHODS

We undertook a cross-sectional study between December 2017 and January 2018 at the Faculties of Medicine and Dentistry University of Belgrade, Serbia. The study comprised 558 students of the sixth year of studies, 430 medical students (response rate 92.47%) and 128 dental students (response rate 81.01%). There were more girls in both samples, 62.8% among medical and 64.1% among dental students.

We used an original semi-structured questionnaire designed for this study to determine knowledge, attitude and practice towards MW management. Students were recruited during their classes and participation was voluntary and anonymous. The questionnaire consisted of fourth parts. Some questions were taken from the questionnaires from similar studies, but they were not standardized, as well as ours.

The first part of questionnaire included questions concerning training in MW management, wearing personal protection (mask, gloves, protective glasses), knowledge about post-exposal prophylaxis and the vaccinal status of students (ten questions). The second part of questionnaire

comprised questions on MW regulation, management (segregation, internal collection, packaging, storage, and final disposal) and injuries reporting system (forty-one question). The third part of questionnaire referred to the knowledge about color coding system (forth questions). The fourth part of questionnaire was in the form of five graded Likert's scale (1= "I fully disagree" 2 = "I mainly disagree"; 3 = "I cannot decide"; 4 = "I mainly agree": 5 = I fully agree") on the statements concerning MW management, continuing training and investigations on this topic (seven questions).

Statistical analysis

We performed statistical analysis with a commercial statistical software SPSS for Windows version 25. We set he significance level at 0.05. The distribution of categorical variables was investigated with χ^2 test. To test the significance of differences between the mean values of numeric and ordinal variables we used Student's t-test and Mann Whitney U test, respectively.

Ethical consideration

We performed the study with the permission from the Ethics Committee of the Faculty of Medicine, University of Belgrade.

RESULTS

The distribution of medical and dental students was similar concerning the training in MW management ($\chi^2 = 1.516$; p > 0.05). A majority of students had no training (74.6% at medical and 79.5% at the dental faculty); a small number of them had partial training (18.5% *vs.* 15.6%); those who had the full training were very few (6.9% *vs.* 4.9%).

Dental students use the protective equipment more frequently than medical students (Table 1). Male students are more disciplined in this regard compared to their female colleagues (56.9% *vs.* 53.0%, respectively; $\chi^2 = 6.446$; p = 0.04). However, vaccinal protection against hepatitis B is better among medical students compared to dental students (Table 1).

Knowledge on post-exposal prophylaxis is better among medical students compared to dental students (44.5% *vs.* 13.3%, respectively, $\chi 2 = 66.308$; p<0.001); the availability of post-exposal prophylaxis is also better at medical faculty compared to dental faculty (36.4% *vs.* 14.8%, $\chi 2 = 31.783$; p<0.001).

Around 80% of students at both faculties are aware of the significance of reporting injuries at work. However, dental students are more disciplined in reporting injuries compared to medical students (63.1% *vs.* 52.4%, $\chi 2 = 4.318$; p = 0.038). The responses of students are similar in relation to the treatment of injuries from sharp objects.

A majority of students are not familiar with the legal regulations regarding medical waste management, as well as with the latest provisions from 2016, but most of them know who the responsible person is for managing the waste at their faculty. Concerning waste separation at the faculty there are more dental students who think that their faculty is separating waste compared to medical students (Table 2).

Concerning the primary separation of medical waste, the students' knowledge on infectious waste and used needles was satisfactory and better among dental students compared to medical students. But, the majority of students at both faculties gave incorrect answers related to chemical and pathoanatomic waste, heavy metals, and cytotoxic drugs (Table 3).

Dental students showed more positive attitude towards medical waste management compared to medical students, particularly concerning the continuing training and investigations on this topic (Cronbach's Alpha = 0.778) (Table 4).

DISCUSSION

The majority of students at both faculties came to the end of studies with no training in the management of medical waste. Similar results were obtained in a study among health professional students in India with only 19% of them who were trained in MW management [7]. Similarly, only about 40% of doctors employed in Health Care Centers in Nigeria, passed adequate training on MW [8]. Another study conducted in Jahor showed that 37% of health workers did not pass adequate training in handling and disposal of sharp objects, with a significantly lower incidence of stabbing on sharp objects among those who were trained in MW compared to those who were not [9].

Dentistry students are disciplined in using protective equipment at work (94.5%); however, the majority of them have not been vaccinated against hepatitis B (59.4%). A minor part of dentistry students is familiar with post-exposal prophylaxis (13.3%), and half of them claim that it is not available. Unlike dental students, medical students are more aware of the significance of vaccination against hepatitis B; but, they use protective equipment in lesser degree (42%), and only a third are familiar with post-exposal prophylaxis. Still, the situation in Serbia in this regard is better than in Nigeria [8] and in Tanzania [10] where post-exposal prophylaxis is familiar to a lesser degree both to

dental and medical students (30.0% and 22.5%, respectively). The relevant results are much better in Jahor, where 87% of medical practitioners confirmed the use of personal protective equipment in handling clinical waste [9].

The vaccinal protection of health workers in Serbia is significantly better than in Nigeria where only 18.5% of health workers are adequately vaccinated against hepatitis B [8]. Medical students are more aware of the significance of vaccination against HBV and are more frequently vaccinated compared to dental students (57.7% *vs.* 39.1%). Immunization against HBV is of utmost importance for all health workers [11]. There are countries that fully recognize this fact and have much better vaccination results. For instance, Saudi Arabia has 80% of dental students vaccinated against hepatitis B [12].

Undergraduate healthcare students have a professional practice, that expose them to biological material. A study conducted in São Paulo, Brazil, registered that 48.8% of students' accidents with biological material occurred among dentistry students, 40.6% among medical students, and 6.5% among nursing students [13]. The practice of reporting injuries resulting from improper waste disposal is very poor in Serbia, in spite of the awareness of reporting significance, confirmed in our study among about 80% of students. We show that only a half of medical students and 63% of dentistry students report injuries from MW; still this is better than in India, where the practice of reporting injuries caused by improper waste disposal is very poor across all groups of health professionals in India [14]. In developing countries, medical workers do not report about 40-75% of injuries from sharp objects; this is a major problem especially in the case of HIV viral infection, where post-exposal prophylaxis effective in 80% of cases [15]. Even in developed countries like Poland injury reporting in hospitals is low; the implementation of regulations in this field did not help [16].

According to the data of the Public Health Institute of Serbia, stab injuries from MW are poorly reported in Serbia; training of health workers may be an effective strategy for improving the practice and behavior towards hospital MW management [17]. In Serbia, 172 seminars were held from 2008 to 2014; 3278 employees in health care were trained at three levels: technician, supervisor and manager of waste management [6]. However, there was no adequate training of students.

A small percentage of Belgrade medical and dental students, 8.2% and 14% respectively, are well informed about the regulations related to MW management. Dental students in India are better informed about MW management; 55.5% of them had good training [18] and 31% of them even know the year when this law was established [19]; in a study conducted in 2016 this number increased to 64.3% [20]. Similarly, to our results, a majority of health workers in Brazil are not familiar with regulations related to MW [21].

Regardless of poor knowledge on regulations, 90% of the investigated students know the responsible person for MW; this is better than in Jahor, where 83% of the respondents know the answer [9]. A recent study showed that the presence of waste managers can effectively minimize the risk of infection [22].

Most of our respondents, especially among dentistry students, know how to dispose of infectious waste and sharp objects; this is similar to the results of a study among the medical staff in the Babol City Hospital, Iran; 97% of them knew MW color coding [23]. But, the majority of students in our study did not know the proper way of managing chemical (90%) or cytotoxic waste (70%). These results are in contrast to a study among dental students from India, where 67% of the participants had a good knowledge about disposing the pharmaceuticals [24]. Similar results were found in a study conducted in Cairo where 60.9% of doctors gave a correct answer regarding the disposal of chemical waste [25]. Better knowledge on the management of infectious waste compared to other hazardous MW is probably related to the dominance of this type of MW in hospitals.

A recent review of the literature has indicated that in many developing countries regulations and laws relating to waste management in hospitals have been adopted recently. However, further knowledge and awareness of the adequate management of medical waste remain poor due to the lack of appropriate training for both doctors and hospital staff [26], although this is a necessity today [27]. A study conducted among Brazilian dental students proved that theoretical knowledge on waste managements can be improved, but this may not amend waste segregation and adequate disposal in dental practice [28]. In our study students show a very positive attitude towards continuing training on MW management and investigations on this topic.

CONCLUSION

In this very first study on knowledge and awareness on MW among Belgrade medical and dental students we show that their training on MW management and vaccinal protection against hepatitis B are unsatisfactory. Dental students show better knowledge on MW and are more disciplined in personal protection compared to medical students. The students support better training on MW management at their faculties and more investigations on this topic.

ACKNOWLEDGMENT

This work has been financially supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, Project No. 175078, 2010-2017.

We would like to express special thanks to Dijana Nikolić, engineer from the Department of Occupational Safety, Faculty of Medicine, University of Belgrade, who helped us in data collection.

REFERENCES

- 1. World Health Organization (WHO) Policy Paper: Safe Health Care Waste Management. Geneva, Switzerland: World Health Organization (WHO); 2014.
- 2. Health care waste. At: http://www.who.int/mediacentre/factsheets/fs253/en/ Accessed: February 22, 2018.
- 3. Milanović S, Jovanovic V, Mršulja A, Stojčić M, Ilić Živojinović J. Medical waste management. A handbook for students of high medical schools. Belgrade: Zavod za udžbenike, 2012. (In Serbian).
- The Law on Waste Management of the Republic of Serbia. Official Gazette, No. 36/2009, 88/2010 and 14/2016
- 5. Jovanović V, Tesink JG, Jovanović D, Đonović N, Jevtić M, Matić B. et al. Development of Healthcare Waste Management in Serbia and Challenges in the Improvement of the Quality of Healthcare Services. Tehnika Kvalitet, standardizacija i metrologija 2014;14(2):343-8.
- 6. Knezevic S, Matic B. A report on the follow-up process on infectious medical waste management within the network of state health facilities in the Republic of Serbia in 2013. Regulated waste flows. Centre for Hygiene and Human Ecology. Republic Institute of Public Health of Serbia. (In Serbian; available at:

http://www.batut.org.rs/download/izvestaji/higijena/2013MedOtpad.pdf).

- Kumar M, Kushwaha R, Kumari M, Singh G, Kumari R. Knowledge, awareness, attitude regarding biomedical waste management among medical students in a tertiary care centre: A cross sectional Study. PARIPEX Indian Journal of Research 2017;6(4):611-4.
- Anozie OB, Lawani LO, Eze JN, Mamah EJ, Onoh RC, Ogah EO et al. Knowledge, Attitude and Practice of Healthcare Managers to Medical Waste Management and Occupational Safety Practices: Findings from Southeast Nigeria. J Clin Diagn Res 2017;11(3):IC01-IC04 doi: 10.7860/JCDR/2017/24230.9527 PMID: 28511409
- Nazli SN, Karuppannan SA/L, Omar D. Knowledge and Awareness of Clinical Waste Management among Medical Practitioners in Hospital Batu Pahat, Johor. IJIMT 2014; 5(2):139-42. doi:10.7763/IJIMT.2014.V5.502
- Mponela MJ, Oleribe OO, Abade A, Kwesigabo G. Post exposure prophylaxis following occupational exposure to HIV: a survey of health care workers in Mbeya, Tanzania, 2009-2010. Pan Afr Med J 2015;21:32. doi: 10.11604/pamj.2015.21.32.4996 PMID: 26405468
- Gupta S, Rani S, Garg S. Infection control knowledge and practice: A cross-sectional survey on dental laboratories in dental institutes of North India. J Indian Prosthodont Soc 2017;17:348-54. doi: 10.4103/jips.jips_5_17. PMID:29249878
- 12. Ahmad IA, Rehan EA, Pani SC. Compliance of Saudi dental students with infection control guidelines. Int Dent J 2013;63(4):196-201. doi: 10.1111/idj.12030. PMID:23879255
- 13. Gir E, Netto JC, Malaguti SE, Canini SRMS, Hayashida M, Machado AA. Accidents with biological material and immunization against hepatitis B among students from the health area. Rev Lat Am Enfermagem 2008;16(3):401-6. doi:10.1590/S0104-11692008000300011
- Vanesh Mathur, S Dwivedi, MA Hassan, and RP Misra. Knowledge, Attitude, and Practices about Biomedical Waste Management among Healthcare Personnel: A Cross-sectional Study. Indian J Community Med 2011;36(2):143–45. doi: 10.4103/0970-0218.84135 PMID: 21976801
- 15. Wilburn SQ. Needlestick and sharps injury prevention. Online J Issues Nurs 2004; 9:5. PMID:15482091
- 16. Garus-Pakowska A, Górajski M, Szatko F. Did legal regulations change the reporting frequency of sharp injuries of medical personnel? Study from 36 hospitals in Łódź Province,

Poland. Int J Occup Med Environ Health 2018;31(1):37-46. doi: 10.13075/ijomeh.1896.01045. PMID:28832030

- Kumar R, Somrongthong R, Ahmed. Impact of waste management training intervention on knowledge, attitude and practices of teaching hospital workers in Pakistan. Pak J Med Sci 2016;32(3):705–710. doi: 10.12669/pjms.323.9903 PMID: 27375718
- Ranjan R, Pathak R, Singh DK, Jalaluddin Md, Kore SA, Kore AR. Awareness about biomedical waste management and knowledge of effective recycling of dental materials among dental students. J Int Soc Prev Community Dent 2016;6(5): 474–479. doi: 10.4103/2231-0762.192941 PMID: 27891315
- Saini R, Pithon MM, Singh HK, Popoff DV. Knowledge of Biomedical Waste Management among the Students of Rural Dental College, Maharashtra, India. Int J Experiment Dent Sci 2013;2:24–6. doi: 10.5005/jp-journals-10029-1034
- Singh LD, Elangbam V, Napolean T, Christina S. Knowledge on Biomedical Waste Management among Medical Students in RIMS, Imphal, Manipur. IOSR-JDMS 2018;17(01):40-43. doi: 10.9790/0853-1701074043
- Assis MC, Gomes VAP, Balista WC, Freitas RR. Use of performance indicators to assess the solid waste management of health services. An Acad Bras Cien 2017;89(3):2445-2460. doi: 10.1590/0001-3765201720170178
- 22. Maina SM, Andrew NK, Caroline WN. Assessment of Level of Knowledge in Medical Waste Management in Selected Hospitals in Kenya. Appli Micro Open Access 2016;2:124. doi: 10.4172/2471-9315.1000124
- Amouei A, Fallah SH, Asgharnia HA, Gholami M, Jafarian S. Knowledge and Attitude of Hospital Personnel Regarding Medical Waste Management. Int Arch Health Sci 2015;2:75-81.
- 24. Sood AG, Sood A. Dental perspective on biomedical waste and mercury management: A knowledge, attitude and practice survey. Indian J Dent Res 2011;22:371-5. doi: 10.4103/0970-9290.87055. PMID:22048573
- Hakim HA, Mohsen A, Bakr I. Knowledge, attitudes and practices of health-care personnel towards waste disposal management at Ain Shams University Hospitals, Cairo. East Mediterr Health J 2014;20(5):347-54. PMID:24952293
- 26. Ali M, Wang W, Chaudhry N, Geng Y. Hospital waste management in developing countries: A mini review. Waste Manag Res 2017;35(6):581-92. Ali M, Wang W, Chaudhry N, Geng Y. Hospital waste management in developing countries: A mini review. Waste Manag Res 2017;35(6):581-92. doi: 10.1177/0734242X17691344 PMID:28566033
- 27. Ozder A, Taker B, Eker HH, Altindis S, Kocaakman M, Karabay O. Medical waste management training for healthcare managers a necessity? J Environ Health Sci Eng 2013;11(1):20. doi:10.1186/2052-336X-11-20 PMID: 24499642
- Victorelli G, Flório FM, Ramacciato JC, Motta RHL, de Souza Fonseca Silva A. Impact of Pedagogical Method on Brazilian Dental Students' Waste Management Practice. J Dent Educ 2014;78(11):1528-33. PMID:25362694

Infection		Facult	у		
Protection	Answer	Medicine n (%)	Dentistry n (%)	Total n (%)	p*
Personal protective devices	No	101 (24.4)	3 (2.3)	104 (19.2)	
	Yes fully	174 (42.0)	121 (94.5)	295 (54.4)	< 0.001
	Yes partially	139 (33.6)	4 (3.1)	143 (26.4)	
	Total	414 (100)	128 (100)	542 (100)	
Vacination	No	159 (37.1)	76 (59.4)	535 (42.3)	< 0.001
Vaccination against hepatitis B	Yes fully	247 (57.7)	50 (39.1)	297 (53.4)	
	Yes partially	22 (5.1)	2 (1.6)	24 (4.3)	
	Total	428 (100)	128 (100)	556 (100)	

Table 1. Infection protection among Belgrade medical and dental students

* χ^2 test

Table 2. Knowledge and attitudes on medical waste management among Belgrade medical	
and dental students	

Knowledge and attitudes on	Answer	Fac	p *		
medical waste management		Medicine n (%)	Dentistry n (%)	F	
Lagal regulation	No	394 (91.8)	111 (86.0)	0.049	
Legal regulation	Yes	35 (8.2)	18 (14.0)	0.049	
Responsible person at the	No	365 (90.3)	119 (93.7)	> 0.05	
faculty	Yes	39 (9.7)	8 (6.3)	> 0.05	
Waste separation at the	No	164 (41.2)	22 (17.2)	.0.001	
faculty	Yes	234 (58.8)	106 (82.8)	< 0.001	
Containers for medical waste	No	55 (13.0)	20 (15.5)	- > 0.05	
at the faculty	Yes	367 (87)	109 (84.5)	> 0.05	

	Faculty				
Primary separation of medical waste	Medicine Answer n (%)/		Den Ans n (p *	
	Correct	Incorrect	Correct	Incorrect	
Infectious waste – yellow	330 (77.8)	94 (22.2)	120 (93.0)	9 (7.0)	< 0.001
Chemical waste – purple	146 (35.4)	266 (64.6)	43 (35.5)	78 (64.5)	> 0.05
Pathoanatomic waste - brown	179 (46.3)	208 (53.7)	50 (43.5)	65 (56.5)	> 0.05
Used needle – yellow	280 (70.4)	118 (29.6)	98 (80.3)	24 (19.7)	0.007
Thermometer with mercury – violet	26 (6.5)	372 (93.5)	12 (9.8)	110 (90.2)	0.012
Syringe – cytotoxic drugs - red	127 (32.7)	261 (67.3)	31 (26.5)	86 (73.5)	> 0.05

Table 3. Knowledge on primary separation of medical waste among Belgrade medical and dental students

 $*\chi^2$ test

Table 4. Attitudes towards medical waste management among Belgrade medical and dental
students (5-graded scale of agreement; mean \pm standard deviation)

Attitude	Medical students	Dental students	p *
Inadequate medical waste management affects human health and environment	$4.72\pm\!\!0.69$	4.71 ± 0.79	0.646
Medical waste management should be a mandatory part of practical and theoretical training of medical and dental students	4.01 ±1.06	4.37 ± 0.92	< 0.001
Training on medical waste management should be performed at least once a year	3.71±1.10	3.96 ± 1.08	0.020
In my opinion medical waste is a topic that deserves more attention in the future	4.27± 0.89	4.49 ± 0.79	0.004
More investigations on medical waste management are needed	3.95 ± 1.12	4.24 ± 1.03	0.005

*Mann–Whitney U-test