



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Tatjana Savić-Stanković, Branislav Karadžić, Marina Latković, Vesna Miletic[†],

**Clinical efficiency of a sodium perborate - hydrogen peroxide mixture
for intracoronar non-vital teeth bleaching**

Клиничка ефикасност микстуре натријум пербората и водоник пероксида
за интракоронарно избељивање депулписаних зуба

University of Belgrade, School of Dental Medicine, Department of Restorative Dentistry and Endodontics,

Belgrade, Serbia

Received: May 4, 2019

Revised: July 24, 2019

Accepted: August 5, 2019

Online First: August 29, 2019

DOI: <https://doi.org/10.2298/SARH190504094S>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. Srp Arh Celok Lek. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

[†]**Correspondence to:**

Vesna MILETIĆ

School of Dental Medicine

Dept. of Restorative Dentistry and Endodontics

Rankeova 4, 11000 Belgrade, Serbia

Email: vesna.miletic@stomf.bg.ac.rs

Clinical efficiency of a sodium perborate - hydrogen peroxide mixture for intracoronary non-vital teeth bleaching

Клиничка ефикасност микстуре натријум пербората и водоник пероксида за интракоронарно избељивање депулписаних зуба

SUMMARY

Introduction/Objective To evaluate initial efficiency of sodium perborate (tetrahydrate) and 30% hydrogen peroxide mixture for intracoronary non-vital teeth bleaching (“walking bleach” technique).

Methods Forty patients with discolored teeth were included in the study. Based on their history and clinical examination, causes of discoloration were classified as necrotic pulp, “endo-sealer” or unknown. The “walking bleach” technique was performed by applying sodium perborate (tetrahydrate) and 30% hydrogen peroxide mixture intracoronally to cavity dentin walls. The mixture was renewed in seven-day intervals. Tooth color was assessed visually before, during, and after the procedure using the Vita Classical shade guide (Vita Zahnfabrik, Bad Säckingen, Germany). Numerical values or shade guide units (SGU) were assigned to Vita shade tabs on a bright-dark scale. Analysis of variance, t-test, correlation and regression analysis were used to analyze the data ($p < 0.05$).

Results On average, 26 ± 9 days or 3–4 appointments were required for intracoronary bleaching to achieve the desired or best possible shade. Better clinical efficiency was found in the necrotic pulp group (17 ± 6 days; 8 ± 3 SGU) than in the “endo-sealer” group (42 ± 13 days; 4 ± 2 SGU) ($p < 0.05$). Age significantly influenced bleaching efficiency ($p < 0.05$). There was no significant correlation between bleaching efficiency and initial shade ($p > 0.05$).

Conclusion Intracoronary, non-vital teeth bleaching (“walking bleach” technique) using sodium perborate (tetrahydrate) and 30% hydrogen peroxide mixture showed satisfactory clinical efficiency. Discoloration caused by pulp necrosis was treated more efficiently than that caused by endodontic sealers. Patient's younger age had a positive effect and discoloration intensity had no effect on bleaching efficiency.

Keywords: walking bleach, intracoronary bleaching, non-vital teeth bleaching, tooth discoloration, sodium perborate, hydrogen peroxide

САЖЕТАК

Увод/циљ Испитати иницијалну клиничку ефикасност микстуре натријумперборат (тетрахидрата) и 30% водоникпероксида за интракоронарно избељивање депулписаних зуба („шетајућа“ техника).

Метод Четрдесет пацијената са пребојеним зубима је било укључено у студију. На бази анамнезе и клиничког прегледа, узроци дисколорације су класификовани као “некротична пулпа”, „ендо-силер“ или „непознат“. Микстура натријум пербората и 30% водоник пероксида апликована је интракоронарно на дентинске зидове кавитета у интервалима од седам дана. Боја зуба је процењивана визуелном методом пре, током и након третмана применом *Vita Classic* кључа боја (*Vita Zahnfabrik*, Немачка). Нумеричка вредност (јединица нијансе) додељена је свакој нијанси *Vita* кључа помоћу светло-тамне скале. Анализа варијансе, *t*-тест, корелациона и регресиона анализа су коришћене за статистичку обраду података ($p < 0,05$).

Резултати Просечно 26 ± 9 дана или 3–4 посете су биле неопходне да се интракоронарним избељивањем постигну жељена или најсветлија могућа нијанса. Боља клиничка ефикасност избељивања забележена је у групи „некротична пулпа“ (17 ± 6 дана; 8 ± 3 нијанси) него у групи „ендо-силер“ (42 ± 13 дана; 4 ± 2 нијансе) ($p < 0,05$). Узраст пацијента је значајно утицао на ефикасност избељивања ($p < 0,05$). Није утврђена значајна повезаност између ефикасности избељивања и почетне нијансе ($p > 0,05$).

Закључак „Шетајућа“ техника избељивања депулписаних зуба применом микстуре натријум перборат (тетрахидрата) и 30% водоник пероксида показала је задовољавајућу клиничку ефикасност. Дисколорација услед некрозе пулпе је успешније избељена него она изазвана ендодонтским пастама. Ефикасније избељивање је забележено код млађих пацијената, док интензитет почетне дисколорације није значајно утицао на ефикасност избељивања.

Кључне речи: шетајућа техника, интракоронарно избељивање зуба, избељивање депулписаних зуба, дисколорација зуба, натријум перборат, водоник пероксид

INTRODUCTION

Tooth color affects smile esthetics, which is among the first facial characteristics observed in a social interaction. Even mild tooth discoloration may reduce individual's self-esteem and quality of life [1]. A recent study showed that a single discolored tooth adversely affects patient's dental self-confidence as well as social and psychological aspects of one's well-being [2].

Non-vital teeth are discolored mainly due to trauma and intracoronary hemorrhage, proteolytic degradation, root canal sealers and cements [3]. Increasingly popular calcium-silicate cements, such as mineral trioxide aggregate (MTA) and Biodentine®, have recently shown to have tooth discoloration potential [4]. Higher discoloration potential has been reported for calcium-silicate-based cements containing bismuth oxide as the radio-pacifying agent and lesser for cements with zirconium oxide [4, 5].

A non-vital tooth bleaching is clinically performed by placing a bleaching agent intracoronally and sealing the crown temporarily. This so-called "walking bleach" technique may be performed in one or more consecutive visits in intervals of three to seven days until the desired or maximum color change is achieved [3]. Regarding color assessment, it should be noted that visual shade matching using Vita shade guides is still most commonly used in daily practice [6].

Sodium perborate, hydrogen peroxide, and carbamide peroxide are commonly used bleaching agents for intracoronary bleaching using the "walking bleach" technique [7, 8]. The efficiency of the bleaching agent may be improved by additional activation with LED blue or laser infrared light or photon-initiated photoacoustic streaming [9, 10]. The mechanism of bleaching by all three agents is based on the release of nascent oxygen and hydrogen peroxide which forms free radicals in the alkaline solution, resulting in hydroperoxyl (HO_2)

and hydroxyl (HO⁻) ions capable of breaking longer-chained chromophores into shorter-chained colorless compounds [11].

In vitro studies of intracoronary bleaching focused on tooth fracture resistance, composition of calcium-silicate cements, their compressive strength and bond strength to composite resin, adhesive bonding, orthodontic brackets bonding to enamel and marginal integrity of temporary and permanent restorative materials following various intracoronary bleaching protocol [12–16].

A similar number of recent clinical studies were found in the literature for in-office bleaching of vital teeth and intracoronary “walking bleach” approach, illustrating the relevance of tooth bleaching in current dental practice and research [1, 2, 7, 8, 9, 17–20]. Hydrogen peroxide, carbamide peroxide and sodium perborate were all-efficient in intracoronary bleaching, with stable effects up to six months or one-year post-bleaching [7, 19, 20]. Even after 25 years post-bleaching 85% of the endodontically treated teeth bleached with 10% carbamide peroxide retained tooth color within two color shades [8]. Multiple studies have confirmed a strong positive effect of both in-office and “walking bleach” procedures on patients’ quality of life, psychosocial interaction, and esthetic self-perception [2, 21]. Regarding the “walking bleach” technique, no difference was found between carbamide and hydrogen peroxide bleaching agents in the context of patient’s esthetic perception, self-confidence, and psychological and social impact [2].

The aim of this clinical study was to evaluate initial efficiency of the sodium perborate - 30% hydrogen peroxide mixture for intracoronary non-vital teeth bleaching (“walking bleach” technique). The null hypotheses were:

1. There is no difference in clinical efficiency of intracoronary bleaching between different causes of discoloration;

2. there is no relationship between clinical efficiency of intracoronaral bleaching and patients' age;
3. there is no relationship between clinical efficiency of intracoronaral bleaching and initial shade.

METHODS

Patient selection, examination, and endodontic treatment

The study was approved by the University of Belgrade, School of Dental Medicine Ethics Committee. Inclusion criteria were visibly discolored crown, non-vital or endodontically treated upper central and lateral incisors, previous endodontic treatment performed two or more years earlier. Exclusion criteria were caries on upper central and lateral incisors, inadequate fillings, and progressive periodontal disease.

Forty patients (25 female and 15 male) aged 18–58 participated in the study. Patients' history was taken and clinical examination performed, including pulp sensitivity tests (electric pulp tester DentVit, Novi Sad, Serbia). Retroalveolar X-rays were taken to check the periapical status and, in cases of previous endodontic treatments, to check the quality of root canal obturation, by assessing the length, conicity and homogeneity of the root canal obturation material. The overall success of previous endodontic treatment was based on patients' history and clinical exam confirming no signs and symptoms of periapical disease. Based on patients' history and examination, potential causes of tooth discoloration were classified into three groups:

1. "Necrotic pulp" (teeth with necrotic pulp with or without patients' recollection of trauma);

2. “endo-sealer” (previously endodontically treated teeth with traces of endo-sealer on pulp chamber walls, patients’ recollection that discoloration occurred post-treatment);
3. “unknown” (previously endodontically treated teeth with no traces of “endo-sealer” on pulp chamber walls, patient could not confirm whether discoloration occurred before or after endodontic treatment).

Initial endodontic treatment was performed in the necrotic pulp group. A re-treatment was performed in previously endodontically treated teeth with inadequate root canal obturation whereas in endodontically treated teeth with adequate root canal obturation the intracoronal bleaching was performed without prior endodontic re-treatment.

Access cavity was prepared and the working length determined using an apex locator (Root ZX, Morita Corp., Osaka, Japan). Root canal instrumentation was performed with Gates-Glidden and K-Flex hand files (Kerr) following the crown-down and step-back technique with 1% sodium hypochlorite irrigation (Patenting, Belgrade, Serbia) Depending on the initial apical size, the master apical files were K#25-35, mostly K#30. Canals were dried with sterile paper points and obturated with gutta-percha (Spident, Korea) and AH-26 sealer (Dentsply, DeTrey, GmbH, Konstanz, Germany) using cold lateral compaction technique. Each tooth was prepared for bleaching by removing 1–2 mm of the obturation material at the canal orifice and sealing it with roughly two mm thick glass-ionomer cement (Vitrebond, 3M ESPE, St. Paul, MN, USA).

Intracoronal bleaching procedure

Tooth color assessment was performed by shade matching with color tabs on the Vita Classical shade guide (Vita Classic, Lumin Vacuum Shade Guide, Vita Zahnfabrik Germany). The procedure was performed by three specialists at daytime in a dental office

with no extra dental unit light. Color assessment was determined by consensus agreement of at least two specialists.

A mixture of sodium perborate (tetrahydrate) and 30% hydrogen peroxide (Sani-Hem, Novi Becej, Srbija) was prepared in the 1:2 ratio by measuring the appropriate quantities of each component on an analytical balance ($d = 0.1$ mg, Acculab, Sartorius, Goettingen, Germany) and mixing them with a metal spatula on a glass slab until the mixture was homogeneous. A portion of the mixture was applied intracoronally as a thin layer and adapted to the entire cavity dentin using a dry, sterile cotton pellet. The cavity was then temporarily sealed with calcium-sulfate cement (Citodur, Dorident, Wien, Austria). At a follow-up after seven days, color assessment was repeated. Temporary filling and the sodium perborate-hydrogen peroxide mixture were removed by an excavator and the cavity was thoroughly rinsed with air-water spray. Freshly prepared mixture was then applied to the inner cavity walls and the tooth was again temporarily restored.

The treatment ended when the desired color was achieved *i.e.* visually acceptable shade in relation to the adjacent teeth or when the color remained unchanged at two consecutive visits. The final achieved shade on the Vital Classical shade guide was recorded as well as the number of required visits.

Following the bleaching procedure, a calcium hydroxide paste was applied to the cavity dentin walls to neutralize the pH. The final restoration was done after two weeks using a two-step total-etch adhesive (Optibond Solo Plus, Kerr, Scafati, Italy) and a micro hybrid composite (XRV Herculite, Kerr, Scafati, Italy). The adhesive was light-cured for 20 s and composite for 40 s using a conventional LED light-curing unit (LEdition, Ivoclar Vivadent, Schaan, Liechtenstein).

All 16 Vita Classical shade tabs were converted to a numerical 1–16 scale, #1 corresponding to the brightest shade (B1) and #16 corresponding to the darkest shade (C4)

(Table 1). For each patient, the start and end point were identified on the numerical scale as shade guide units. Intracoronal bleaching efficiency (BE) was expressed as the percentage of the bleached shade relative to the initial shade on the bright-dark scale.

$$\text{BE (\%)} = [(\text{Initial shade} - \text{End shade}) / \text{Initial shade}] \times 100$$

Statistical analysis

Data were statistically analyzed using analysis of variance, Student's t test, correlation and regression analyses at the level of significance $\alpha = 0.05$. Statistical analysis was performed in Minitab 16 (Minitab Inc. State College, PA, USA).

RESULTS

The age structure *i.e.* mean (standard deviation) of 15 male and 25 female patients involved in this study was comparable: 37 (10.4) years of male and 32.6 (10.6) years of female patients. Necrotic pulp was diagnosed in 16 patients, 14 patients were allocated to the “endo-sealer” group, and in 10 patients, the cause of discoloration was unknown. All data taken together, 26 ± 9 days or 3–4 appointments were required for the intracoronal bleaching procedure to achieve the desired or best possible shade.

Table 2 presents information on the treatment duration relative to the cause of discoloration. Significantly shorter treatment was required in the necrotic pulp group than in the “endo-sealer” group ($p < 0.05$). In addition to longer mean treatment time, the “endo-sealer” group showed larger variation in terms of minimum and maximum duration spanning to almost 50 days compared to the necrotic pulp group with only 18 days between the minimum and maximum treatment duration. No significant differences were found between the previous two groups and the unknown group ($p > 0.05$). The number of patients in each group indicates that all patients returned for follow-ups *i.e.* the recall rate was 100%.

All data taken together, color change of 7 ± 2 shade guide units could be achieved. Table 3 presents data relative to the cause of discoloration. There was a statistically significant difference between the number of shade guide units achieved by bleaching in the necrotic pulp than in the “endo-sealer” group ($p < 0.05$). The unknown group was comparable to both previous groups ($p > 0.05$). In all three groups, the minimum number of shade guide units was the same (3), but the necrotic pulp group exceeded the other two groups in the number of maximum achievable shade guide units during bleaching (14).

Figure 1 shows a significant relationship between BE and patients' age. Pearson correlation analysis showed negative correlation between BE and patients' age ($r = -0.4708$; $p = 0.002$), indicating that better clinical efficiency could be accomplished in younger patients. Correlation and regression analyses showed no significant relationship between BE and initial shade ($r = -0.2826$; $p = 0.077$) (Figure 2).

Figures 3 and 4 show a case of intracoronal bleaching of an upper right central incisor discolored by trauma. Figures 5 and 6 show a case of intracoronal bleaching of an upper left central incisor discolored by “endo-sealer”.

DISCUSSION

The first and second null hypotheses were rejected as different causes of tooth discoloration had a significant effect on intracoronal bleaching efficiency, which was more effective in younger patients. The third null hypothesis was upheld as no significant relationship was established between bleaching efficiency and initial shade.

This clinical study showed the potential of a sodium perborate (tetrahydrate) mixture with 30% of hydrogen peroxide to bleach discolored teeth. On average, the desired or maximum bleaching effect for this “walking bleach” technique required 3–4 appointments and 26 days. The present results are comparable to other recent clinical studies on the

efficiency of the “walking bleach” technique. Pedrollo Lise et al. [7] used a mixture of sodium perborate (unspecified monohydrate, trihydrate, or tetrahydrate) and 20% hydrogen peroxide up to four weeks post-bleaching. Bersezio et al. [22] compared 35% hydrogen peroxide to 37% carbamide peroxide and Gupta et al. [20] used sodium perborate (tetrahydrate) alone for the same purpose. Both studies reported a similar number of required appointments for the most desirable outcome, 3–4 in Bersezio et al. [22] and three in Gupta et al. [20]. Given the fact that in all the studies, different mixtures were applied for intracoronary bleaching, it may be concluded that similar treatment longevity *i.e.* the number of appointments is expected irrespective of the bleaching agent/agents.

The desired or maximum achievable effect of intracoronary bleaching in the present study was seven shades on average. This was fewer than 12 shades reported by Gupta et al. [20] or 15 shades achieved by Bersezio et al. [22]. The differences could be explained by the greater number of teeth discolored by “endo-sealer” reducing the number of achieved shades in the present study. Gupta et al. [20] treated 41 patients with discolored teeth due to trauma, but without previous endodontic treatment, meaning that not a single “endo-sealer” case was included in their study. Nearly half of the 47 treated patients in Bersezio et al. [22] had discoloration due to trauma, whilst the other half of discolored teeth were previously endodontically treated but “endo-sealer” was not the stated cause of discoloration.

Generally, the necrotic pulp group showed a higher efficiency of intracoronary bleaching in the present study than the “endo-sealer” group both in terms of the required time/number of appointments and of the shade difference in bleached teeth. Despite the fact that the cause of discoloration was based on patients’ history and clinical examination and not confirmed by chemical testing, the necrotic pulp and “endo-sealer” causes could be differentiated with high confidence. The same approach was followed in previously mentioned studies [7, 20, 22]. Furthermore, a distinct difference in the efficiency of bleaching confirms the correct

diagnosis of the cause of discoloration. The unknown group showed the results between the other two groups indicating that it probably contained teeth with both causes of discoloration. The observed differences between the necrotic pulp and “endo-sealer” groups could be associated with the fact that pigments of natural origin (decomposition products of blood and pulp tissue components as well as pigments of microbial origin) are more susceptible to oxidation reactions that lead to chromophore cleavage and molecular conversion than endodontic sealers [23]. Endodontic sealers containing silver, bismuth and resins cause not only discernible discoloration shortly after placement but also show a significant degree of bleaching resistance [24]. Discoloration induced by endodontic sealers may be more bleaching resistant than that induced by natural pigments due to the amount of excess sealer in the pulp chamber and complex interactions between sealer components and dentin tissue [25]. For example, it was recently explained that bismuth from MTA-based materials forms black precipitates upon interaction with dentin collagen in the presence of sodium hypochlorite, bismuth carbonate upon reaction with atmospheric carbon dioxide or breaks down to metallic bismuth and oxygen upon light irradiation in oxygen-free environment [26, 27].

Better treatment efficiency was associated with patients' younger age and no direct relationship was found between bleaching efficiency and initial shade. These results are comparable to Gupta et al. [20] who also performed intracoronary non-vital teeth bleaching. Based on pooled data from 11 clinical trials, patients' younger age was reported as a significant factor in clinical efficiency of vital teeth bleaching as well [28]. More efficient bleaching in younger patients and higher bleaching resistance in older patients may be associated with the diameter of dentinal tubules. In younger patients, wider dentinal tubules allow deeper penetration of hydroperoxyl (HO_2) and hydroxyl (HO^-) ions. Conversely, the results indicated that even a highly discolored tooth with necrotic pulp could be efficiently

bleached, albeit taking more time and appointments, than a tooth mildly discolored by endodontic sealers. The cause was shown to have a more important effect on treatment efficiency than the intensity of initial discoloration.

It could be argued that in-office non-vital bleaching is faster than the “walking bleach” technique. Although this is true, the “walking bleach” technique is a more conservative approach that allows better control of the achieved result over a longer time. More aggressive in-office non-vital bleaching could lead to over-bleaching due to limited color control. Furthermore, studies indicate that the “walking bleach” technique results in a stable color over a period of months or even years [7, 8, 19]. The cost of commercial products for in-office bleaching often increases the cost of this treatment. Conversely, the low cost of sodium perborate powder and hydrogen peroxide used in the “walking bleach” technique makes this technique financially acceptable.

Despite advantages, biological risks of the “walking bleach” technique were recently investigated by Bersezio et al. [29]. They reported increase in the production of IL-1 β and RANKL biomarkers in periodontal tissues by analyzing gingival crevicular fluid up to six months after intracoronal bleaching with either 35% hydrogen peroxide or 37% carbamide peroxide over four sessions. Increased production of these biomarkers may be associated with osteoclastic activity with a potential for root resorption as an adverse consequence of intracoronal bleaching [30, 31].

A limitation of this study is that the efficiency of intracoronal bleaching was assessed visually using a Vita Classical shade guide. However, the same approach was followed in other studies [8, 20, 22]. Though digital shade matching has proven more accurate and reliable than visual shade matching, experience and training may positively affect accuracy of visual shade matching [32, 33]. Colorimeters and spectrophotometers have not been widely accepted probably due to high costs of such devices. The fact that visual shade matching was

performed in the present study ensures clinically relevant results as most dentists would likely resort to the same technique when assessing bleaching efficiency. Furthermore, the 50:50 perceptibility and acceptability thresholds in color observation in dentistry was found to be significantly lower for dentists (mean ΔE_{00} 0.62 and 1.79, respectively) than laypersons (mean ΔE_{00} 1.00 and 2.04, respectively) who largely constitute the patient group [33]. This indicates that dentists observe color differences better than patients, likely due to training, ensuring better treatment results and patient satisfaction.

CONCLUSION

Intracoronary, non-vital teeth bleaching or the “walking bleach” technique using a mixture of sodium perborate tetrahydrate and 30% hydrogen peroxide showed satisfactory clinical efficiency. On average, 3–4 appointments were required for a desired or maximum possible outcome. The cause of discoloration affected treatment longevity and efficiency, in that discoloration due to pulp necrosis was resolved more efficiently than discoloration due to endodontic sealers. Patients’ younger age had a positive effect and discoloration intensity had no effect on bleaching efficiency.

Conflict of interest: None declared.

REFERENCES

1. Pavicic DK, Kolceg M, Lajnert V et al. Changes in Quality of Life Induced by Tooth Whitening are Moderated by Perfectionism: A Randomized, Double-Blind, Placebo-Controlled Trial. *Int J Prosthodont*. 2018;31:394-6. doi: 10.11607/ijp.5499. PMID: 29624630
2. Bersezio C, Martin J, Mayer C et al. Quality of life and stability of tooth color change at three months after dental bleaching. *Qual Life Res*. 2018;27:3199-207. doi: 10.1007/s11136-018-1972-7. PMID:
3. Plotino G, Buono L, Grande NM et al. Nonvital tooth bleaching: a review of the literature and clinical procedures. *J Endod*. 2008;34:394-407. doi: 10.1016/j.joen.2007.12.020. PMID: 18358884
4. Mozynska J, Metlerski M, Lipski M, Nowicka A. Tooth Discoloration Induced by Different Calcium Silicate-based Cements: A Systematic Review of In Vitro Studies. *J Endod*. 2017;43:1593-601. doi: 10.1016/j.joen.2017.04.002. PMID: 28864217
5. Marconyak LJ, Jr., Kirkpatrick TC, Roberts HW et al. A Comparison of Coronal Tooth Discoloration Elicited by Various Endodontic Reparative Materials. *J Endod*. 2016;42:470-3. doi: 10.1016/j.joen.2015.10.013. PMID: 26620853
6. Wee AG, Meyer A, Wu W, Wichman CS. Lighting conditions used during visual shade matching in private dental offices. *J Prosthet Dent*. 2016;115:469-74. doi: 10.1016/j.prosdent.2015.09.020. PMID: 26723088
7. Pedrollo Lise D, Siedschlag G, Bernardon JK, Baratieri LN. Randomized clinical trial of 2 nonvital tooth bleaching techniques: A 1-year follow-up. *J Prosthet Dent*. 2018;119:53-9. doi: 10.1016/j.prosdent.2017.03.004. PMID: 28478984
8. Amato A, Caggiano M, Pantaleo G, Amato M. In-office and walking bleach dental treatments on endodontically-treated teeth: 25 years follow-up. *Minerva Stomatol*. 2018;67:225-30. doi: 10.23736/S0026-4970.18.04190-0. PMID: 30499284
9. Vildosola P, Bottner J, Avalos F et al. Teeth bleaching with low concentrations of hydrogen peroxide (6%) and catalyzed by LED blue (450 +/- 10 nm) and laser infrared (808 +/- 10 nm) light for in-office treatment: Randomized clinical trial 1-year follow-up. *J Esthet Restor Dent*. 2017;29:339-45. doi: 10.1111/jerd.12318. PMID: 28712152
10. Arslan H, Akcay M, Yasa B et al. Bleaching effect of activation of hydrogen peroxide using photon-initiated photoacoustic streaming technique. *Clin Oral Investig*. 2015;19:253-9. doi: 10.1007/s00784-014-1255-9. PMID: 24859380
11. Joiner A. The bleaching of teeth: a review of the literature. *J Dent*. 2006;34:412-9. doi: 10.1016/j.jdent.2006.02.002. PMID: 16569473
12. Oskoe SS, Bahari M, Daneshpooy M et al. Effect of Different Intraorifice Barriers and Bleaching Agents on the Fracture Resistance of Endodontically Treated Anterior Teeth. *J Endod*. 2018;44:1731-5. doi: 10.1016/j.joen.2018.07.025. PMID: 30266467
13. Keskin C, Sariyilmaz E, Kele SA. The effect of bleaching agents on the compressive strength of calcium silicate-based materials. *Aust Endod J*. 2018doi: 10.1111/aej.12318. PMID: 30338618
14. Cavalli V, Sebold M, Shinohara MS et al. Dentin bond strength and nanoleakage of the adhesive interface after intracoronary bleaching. *Microsc Res Tech*. 2018;81:428-36. doi: 10.1002/jemt.22995. PMID: 29345388
15. Chauhan V, Kumar P, Sharma P, Shetty D. Effect of different intracoronary bleaching methods on shear bond strength of ceramic brackets bonded to bleached enamel: An in-vitro study. *J Orthod Sci*. 2017;6:86-90. doi: 10.4103/jos.JOS_149_16. PMID: 28717632
16. Srikumar GP, Varma KR, Shetty KH, Kumar P. Coronal microleakage with five different temporary restorative materials following walking bleach technique: An ex-vivo study. *Contemp Clin Dent*. 2012;3:421-6. doi: 10.4103/0976-237X.107431. PMID: 23633802
17. Martins I, Onofre S, Franco N et al. Effectiveness of In-office Hydrogen Peroxide With Two Different Protocols: A Two-center Randomized Clinical Trial. *Oper Dent*. 2018;43:353-61. doi: 10.2341/17-128-C. PMID: 29949479
18. Klaric Sever E, Budimir Z, Cerovac M et al. Clinical and patient reported outcomes of bleaching effectiveness. *Acta Odontol Scand*. 2018;76:30-8. doi: 10.1080/00016357.2017.1376111. PMID: 28893130

19. Bersezio C, Ledezma P, Mayer C et al. Effectiveness and effect of non-vital bleaching on the quality of life of patients up to 6 months post-treatment: a randomized clinical trial. *Clin Oral Investig*. 2018;22:3013-9. doi: 10.1007/s00784-018-2389-y. PMID: 29455372
20. Gupta SK, Saxena P. Evaluation of patient satisfaction after non-vital bleaching in traumatized discolored intact anterior teeth. *Dent Traumatol*. 2014;30:396-9. doi: 10.1111/edt.12109. PMID: 24720860
21. Bersezio C, Estay J, Jorquera G et al. Effectiveness of Dental Bleaching With 37.5% and 6% Hydrogen Peroxide and Its Effect on Quality of Life. *Oper Dent*. 2019;44:146-55. doi: 10.2341/17-229-C. PMID: 30517065
22. Bersezio C, Martin J, Pena F et al. Effectiveness and Impact of the Walking Bleach Technique on Esthetic Self-perception and Psychosocial Factors: A Randomized Double-blind Clinical Trial. *Oper Dent*. 2017;42:596-605. doi: 10.2341/16-133-C. PMID: 29144879
23. Watts A, Addy M. Tooth discoloration and staining: a review of the literature. *Br Dent J*. 2001;190:309-16. doi: 10.1038/sj.bdj.4800959a. PMID: 11325156
24. Gurel MA, Kivanc BH, Ekici A, Alacam T. Evaluation of crown discoloration induced by endodontic sealers and colour change ratio determination after bleaching. *Aust Endod J*. 2016;42:119-23. doi: 10.1111/aej.12147. PMID: 27134011
25. Meincke DK, Prado M, Gomes BP et al. Effect of endodontic sealers on tooth color. *J Dent*. 2013;41 Suppl 3:e93-6. doi: 10.1016/j.jdent.2012.10.011. PMID: 23123496
26. Marciano MA, Duarte MA, Camilleri J. Dental discoloration caused by bismuth oxide in MTA in the presence of sodium hypochlorite. *Clin Oral Investig*. 2015;19:2201-9. doi: 10.1007/s00784-015-1466-8. PMID: 25922130
27. Valles M, Roig M, Duran-Sindreu F et al. Color Stability of Teeth Restored with Biodentine: A 6-month In Vitro Study. *J Endod*. 2015;41:1157-60. doi: 10.1016/j.joen.2015.03.014. PMID: 25937179
28. Rezende M, Loguercio AD, Kossatz S, Reis A. Predictive factors on the efficacy and risk/intensity of tooth sensitivity of dental bleaching: A multi regression and logistic analysis. *J Dent*. 2016;45:1-6. doi: 10.1016/j.jdent.2015.11.003. PMID: 26612623
29. Bersezio C, Estay J, Saez M et al. Six-month Follow-up of the Effect of Nonvital Bleaching on IL-1beta and RANK-L: A Randomized Clinical Trial. *Oper Dent*. 2019doi: 10.2341/18-023-C. PMID: 31034350
30. Huynh NC, Everts V, Pavasant P, Ampornaramveth RS. Interleukin-1beta induces human cementoblasts to support osteoclastogenesis. *Int J Oral Sci*. 2017;9:e5. doi: 10.1038/ijos.2017.45. PMID: PMC5729550
31. Kim Y, Lee CY, Kim E, Roh BD. Invasive cervical resorption: treatment challenges. *Restor Dent Endod*. 2012;37:228-31. doi: 10.5395/rde.2012.37.4.228. PMID: PMC3568643
32. Brandt J, Nelson S, Lauer HC et al. In vivo study for tooth colour determination-visual versus digital. *Clin Oral Investig*. 2017;21:2863-71. doi: 10.1007/s00784-017-2088-0. PMID: 28265772
33. Paravina RD, Ghinea R, Herrera LJ et al. Color difference thresholds in dentistry. *J Esthet Restor Dent*. 2015;27 Suppl 1:S1-9. doi: 10.1111/jerd.12149. PMID: 25886208

Table 1. Bright – dark scale for numerical expression of the Vita Classical shade tabs as shade guide units.

Vita tab	B1	A1	B2	D2	A2	C1	C2	D4	A3	D3	B3	A3.5	B4	C3	A4	C4
Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16

Paper accepted

Table 2. Treatment duration (days) in different etiological groups.

Duration in days	Necrotic pulp	Endo-sealer	Unknown
Mean	17	42	29
Standard deviation	6	13	11
Minimum	3	21	14
Maximum	21	70	42
Number of patients	16	14	10

Paper accepted

Table 3. Changes in shade guide units following intracoronal bleaching.

Changes in shade units	Necrotic pulp	Endo-sealer	Unknown
Mean	8	4	6
Standard deviation	3	2	2
Minimum	3	3	3
Maximum	14	11	10
Number of patients	16	14	10

Paper accepted




Figure 1. Regression analysis showing a significant relationship between bleaching efficiency and patients' age.




Figure 2. Regression analysis showing no significant relationship between bleaching efficiency and initial shade.



Figure 3. An upper right central incisor discolored by trauma before intracoronal bleaching.

Paper accepted



Figure 4. The same upper right central incisors discolored by trauma after intracoronal bleaching.

Paper accepted



Figure 5. An upper left incisor discolored by endo-sealer before intracoronal bleaching.

Paper accepted



Figure 6. The same upper left incisor discolored by endo-sealer after intracoronal bleaching.

Paper accepted